arXiv:0710.2166v2 [math.GT] 27 Mar 2008

LOCAL TORUS ACTIONS MODELED ON THE STANDARD REPRESENTATION

TAKAHIKO YOSHIDA

Abstract. We introduce the notion of a *local torus action modeled on the standard representation* (for simplicity, we call it a local torus action). It is a generalization of a locally standard torus action and also an underlying structure of a locally toric Lagrangian fibration. For a local torus action, we define two invariants called a *characteristic pair* and an *Euler class of the orbit map*, and prove that local torus actions are classified topologically by them. As a corollary, we obtain a topological classification of locally standard torus actions, which is a generalization of the topological classification of quasi-toric manifolds by Davis-Januszkiewicz [10, Proposition 1.8] and of effective T^2-actions on four-dimensional manifolds without nontrivial finite stabilizers by Orlik-Raymond [31]. We investigate locally toric Lagrangian fibrations from the viewpoint of local torus actions. We give a necessary and sufficient condition in order that a local torus action becomes a locally toric Lagrangian fibration. Locally toric Lagrangian fibrations are classified by Boucetta-Molino [5] up to fiber-preserving symplectomorphisms. We shall reprove the classification theorem of locally toric Lagrangian fibrations by refining the proof of the classification theorem of local torus actions. We also investigate the topology of a manifold equipped with a local torus action when the Euler class of the orbit map vanishes.

1. Introduction

Let S^1 be the unit circle in $\mathbb{C}$ and $T^n := (S^1)^n$ the n-dimensional compact torus. T^n acts on the n-dimensional complex vector space $\mathbb{C}^n$ by coordinatewise complex multiplication. This action is called the *standard representation of* T^n. In this paper we focus on manifolds which are locally modeled on the standard representation of T^n. A typical example is a nonsingular toric variety. T^n acts on an n-dimensional toric variety as a subgroup of the n-dimensional complex torus $(\mathbb{C}^*)^n$. If the toric variety is nonsingular, then for each point of it we can take a coordinate neighborhood (U, ρ, φ), where U is a T^n-invariant connected open neighborhood of the point, ρ is an automorphism of T^n, and φ is a ρ-equivariantly diffeomorphism from U to an open set of $\mathbb{C}^n$ invariant under the standard representation of T^n. The latter means that $\varphi(u \cdot x) = \rho(u) \cdot \varphi(x)$ for $u \in T^n$ and $x \in U$. In general, a torus action which has an atlas consisting of such coordinate neighborhoods is said to be *locally standard* and such an atlas is called a *standard atlas*. This is one of the starting point of their pioneer work [10] of Davis-Januszkiewicz, in which they focused on locally standard torus actions whose orbit spaces are simple convex polytopes and showed that they still have fascinating combinatorial properties as well as the original toric varieties. (A manifold with this torus action is now called a *quasi-toric manifold*.) After their work, topological generalizations of the original toric theory have been actively studied [6, 7, 21, 22, 23, 26, 27].

2000 *Mathematics Subject Classification.* Primary 57R15; Secondary 57S99, 55R55.

Key words and phrases. local torus actions, locally standard torus actions, Lagrangian fibrations.

The author was partially supported by Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists.

In this paper we consider a structure similar to a standard atlas, but which satisfies a weaker condition than that of a standard atlas.

Definition 1.1. Let X be a compact Hausdorff space. A *weakly standard* C^r $(0 \leq r \leq \infty)$ *atlas* of X is an atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha \in \mathcal{A}}$ which satisfies the following properties

- (i) for each α, φ_α^X is a homeomorphism from U_α^X to an open set of $\mathbb{C}^n$ which is invariant with respect to the standard representation of T^n,
- (ii) on each nonempty overlap $U_{\alpha\beta}^X := U_\alpha^X \cap U_\beta^X$,
  - (a) $\varphi_\alpha^X(U_{\alpha\beta}^X)$ and $\varphi_\beta^X(U_{\alpha\beta}^X)$ are also invariant with respect to the standard representation of T^n and
  - (b) there exists an automorphism $\rho_{\alpha\beta}$ of T^n such that the overlap map $\varphi_{\alpha\beta}^X := \varphi_\alpha^X \circ (\varphi_\beta^X)^{-1}$ is $\rho_{\alpha\beta}$-equivariant C^r diffeomorphic with respect to the standard representation of T^n.[1]

Two weakly standard C^r atlases $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha \in \mathcal{A}}$ and $\{(V_\beta^X, \psi_\beta^X)\}_{\beta \in \mathcal{B}}$ of X are defined to be *equivalent* if on each nonempty overlap $U_\alpha^X \cap V_\beta^X$, there exists an automorphism ρ of T^n such that $\varphi_\alpha^X \circ (\psi_\beta^X)^{-1}$ is ρ-equivariant C^r diffeomorphic. We call an equivalence class of weakly standard C^r atlases a C^r *local* T^n*-action on* X *modeled on the standard representation*, or a local T^n-action on X if there are no confusions and denote it by $\mathcal{T}$.

Definition 1.2. Let $(X_i, \mathcal{T}_i)$ $(i = 1, 2)$ be a $2n$-dimensional manifold equipped with a C^r local T^n-action $\mathcal{T}_i$ and $\{(U_\alpha^{X_1}, \varphi_\alpha^{X_1})\}_{\alpha \in \mathcal{A}} \in \mathcal{T}_1$ and $\{(U_\beta^{X_2}, \varphi_\beta^{X_2})\}_{\beta \in \mathcal{B}} \in \mathcal{T}_2$ the maximal weakly standard atlases of X_1 and X_2. $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ are said to be C^r *isomorphic* if there exists a C^r diffeomorphism $f_X \colon X_1 \to X_2$, and there exists an automorphism ρ of T^n on each nonempty overlap $U_\alpha^{X_1} \cap f_X^{-1}(U_\beta^{X_2}) \neq \emptyset$ such that $\varphi_\beta^{X_2} \circ f_X \circ (\varphi_\alpha^{X_1})^{-1}$ is ρ-equivariant. f_X is called a C^r *isomorphism* and we denote it by $f_X \colon (X_1, \mathcal{T}_1) \to (X_2, \mathcal{T}_2)$.

The purpose of this paper is to classify local torus actions topologically in terms of certain invariants. This is an improvement on the previous work [33]. The main motivation of this work is to develop the equivariant theory of local torus actions and its application to geometry and topology of Lagrangian fibrations, which we are investigating now. In [34] we also discuss the lifting problem of local torus actions to fiber bundles.

It is obvious that a standard atlas of a locally standard torus action satisfies the conditions in Definition 1.1. In this sense, the notion of a local torus action is a generalization of that of a locally standard torus action. As in the usual group action case, we can define the orbit space and the orbit map for a local T^n-action $\mathcal{T}$ on a manifold X. In fact, we construct an n-dimensional C^0 manifold B_X with corners and a C^0 open map $\mu_X \colon X \to B_X$ which is locally identified with the orbit map of the standard representation of T^n in Section 2.

Locally standard torus actions provide several examples of local torus actions. But, not all local torus actions come from locally standard torus actions. Let $\mathrm{Aut}(T^n)$ be the group of automorphisms of T^n as a group. Then, a nontrivial T^n-bundle on an n-dimensional closed manifold whose structure group is $\mathrm{Aut}(T^n)$ is equipped with a local T^n-action which is not induced by any locally standard T^n-action. In general, for a local T^n-action $\mathcal{T}$ on a manifold X, a weakly standard atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha \in \mathcal{A}} \in \mathcal{T}$ induces an atlas $\{(U_\alpha^B, \varphi_\alpha^B)\}$ of B_X, and automorphisms $\rho_{\alpha\beta}$ in (ii) of Definition 1.1 form a Čech one-cocycle on $\{U_\alpha^B\}_{\alpha \in \mathcal{A}}$ with values in $\mathrm{Aut}(T^n)$, hence it defines a cohomology class in the first Čech cohomology of B_X

---

[1] In this paper, for simplicity, we assume that every overlap of two open sets is connected.

with coefficients in $\mathrm{Aut}(T^n)$. We show that it is the obstruction for a local T^n-action to come from a locally standard T^n-action (see Proposition 3.5).

To give a topological classification of local torus actions, let us first recall the topological classification of quasi-toric manifolds by Davis-Januszkiewicz. In [10] Davis-Januszkiewicz defined an invariant called a *characteristic function* for a quasi-toric manifold. They constructed a new quasi-toric manifold, called a canonical model, by using the characteristic function, and proved that a quasi-toric manifold is equivariant homeomorphic to the canonical model of it. Masuda-Panov generalized their technique to locally standard torus manifolds and proved that the same is true for a locally standard torus manifold provided that the second cohomology group of the orbit space vanishes. We shall classify local torus actions topologically by generalizing their method. For a local T^n-action $\mathcal{T}$ on a manifold X and a weakly standard atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$, the Čech one-cocycle $\{\rho_{\alpha\beta}\}$ corresponds to the principal $\mathrm{Aut}(T^n)$-bundle which we denote by $\pi_{P_X} \colon P_X \to B_X$ on B_X. By Proposition 3.5, when $(X, \mathcal{T})$ comes from a locally standard T^n-action, P_X is the product bundle $P_X = B_X \times \mathrm{Aut}(T^n)$. Let Λ be the lattice of integral elements in the Lie algebra of T^n, and we denote by $\pi_{\Lambda_X} \colon \Lambda_X \to B_X$ the associated Λ-bundle of P_X by the natural isomorphism between $\mathrm{Aut}(T^n)$ and the automorphism group $\mathrm{GL}(\Lambda)$ of Λ. Since B_X is a manifold with corners, there is a natural stratification on B_X. We denote by $\mathcal{S}^{(n-1)}B_X$ the codimension one part of the natural stratification of B_X. For $(X, \mathcal{T})$, let $\pi_{T_X} \colon T_X \to B_X$ the associated T^n-bundle of P_X by the natural action of $\mathrm{Aut}(T^n)$ on T^n. By construction, T_X acts fiberwise on X. So a characteristic function is generalized to a rank one subbundle $\pi_{\mathcal{L}_X} \colon \mathcal{L}_X \to \mathcal{S}^{(n-1)}B_X$ of the restriction of $\pi_{\Lambda_X} \colon \Lambda_X \to B_X$ to $\mathcal{S}^{(n-1)}B$. We call $\pi_{\mathcal{L}_X} \colon \mathcal{L}_X \to \mathcal{S}^{(n-1)}B_X$ the *characteristic bundle* and also call the pair $(P_X, \mathcal{L}_X)$ the *characteristic pair*. Similar to quasi-toric manifolds, we can construct a new C^0 manifold $X_{(P_X,\mathcal{L}_X)}$ equipped with a local torus action by using $(P_X, \mathcal{L}_X)$. We still call $X_{(P_X,\mathcal{L}_X)}$ a canonical model. But unlike the above case, in general, $X_{(P_X,\mathcal{L}_X)}$ is not C^0 isomorphic to $(X, \mathcal{T})$ even if $(X, \mathcal{T})$ comes from a locally standard torus action. So, in order to represent the difference between them, we define another topological invariant $e(X, \mathcal{T})$ called the *Euler class of* μ_X. It is a natural generalization of an Euler class of a principal torus bundle. We also show that $e(X, \mathcal{T})$ is an obstruction in order that μ_X admits a C^0 section. With these preliminaries we can state a topological classification theorem of local torus actions (Theorem 6.1).

Theorem 1 (a topological classification of local torus actions)**.** *Local torus actions are classified by the characteristic pairs and the Euler classes of the orbit maps up to C^0 isomorphisms.*

We focus on locally standard torus actions. Then we can also obtain the following corollary (Corollary 6.3).

Corollary 2 (a topological classification of locally standard torus actions)**.** *Locally standard torus actions are classified by the characteristic bundles and the Euler classes of the orbit maps up to equivariant homeomorphisms.*

This is a generalization of the topological classification of quasi-toric manifolds by Davis-Januszkiewicz [10] and of effective T^2-actions on four-dimensional manifolds without nontrivial finite stabilizers by Orlik-Raymond [31].

In case of $e(X, \mathcal{T}) = 0$, we shall investigate the topology of a manifold $(X, \mathcal{T})$ with a local torus action by using algebraic topological methods. We show that if the zero-dimensional part $\mathcal{S}^{(0)}B_X$ of B_X is nonempty, then the fundamental group of X is isomorphic to the one of B_X (Theorem 8.1). We give a way to compute cohomology groups and K-groups by using the Atiyah-Hirzebruch-Leray spectral sequence for μ_X (Theorem 8.2 and Theorem 8.6). In the case where X is four-dimensional and both X and B_X are oriented, we observe the signature of X.

Another important class of local torus actions appears in Lagrangian fibrations. Let $\omega_{\mathbb{C}^n}$ be the standard symplectic structure on $\mathbb{C}^n$

$$\omega_{\mathbb{C}^n} := \frac{1}{2\pi\sqrt{-1}} \sum_{k=1}^{n} dz_k \wedge d\overline{z}_k. \tag{1.1}$$

The standard representation of T^n preserves $\omega_{\mathbb{C}^n}$ and the map $\mu_{\mathbb{C}^n} \colon \mathbb{C}^n \to \mathbb{R}^n$ which is defined by

$$\mu_{\mathbb{C}^n}(z) := (|z_1|^2, \dots, |z_n|^2) \tag{1.2}$$

for $z = (z_1, \dots, z_n) \in \mathbb{C}^n$ is a moment map of the standard representation of T^n. Note that the image of $\mu_{\mathbb{C}^n}$ is the n-dimensional standard positive cone

$$\mathbb{R}^n_+ := \{\xi = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n \colon \xi_i \geq 0 \ i = 1, \dots, n\}. \tag{1.3}$$

Let (X, ω) be a $2n$-dimensional closed C^∞ symplectic manifold and B an n-dimensional C^∞ manifold with corners. A C^∞ map $\mu \colon (X, \omega) \to B$ is called a *locally toric Lagrangian fibration* if it is locally identified with $\mu_{\mathbb{C}^n} \colon (\mathbb{C}^n, \omega_{\mathbb{C}^n}) \to \mathbb{R}^n_+$ (for the precise definition see Definition 3.7). It is a natural generalization of a moment map of a symplectic toric manifold. In the case of $\partial B = \emptyset$, it is a nonsingular Lagrangian fibration. Conversely, by the Arnold-Liouville theorem [1], a nonsingular Lagrangian fibration with closed connected fibers on a closed manifold is also such an example. We will see in Proposition 3.13 that for a locally toric Lagrangian fibration $\mu \colon (X, \omega) \to B$, X admits a C^∞ local torus action. We also give a necessary and sufficient condition in order that a C^∞ local torus action $(X, \mathcal{T})$ admits a symplectic structure ω so that $\mu_X \colon (X, \omega) \to B_X$ is a locally toric Lagrangian fibration (Theorem 7.7). In particular, as is well-known, if $\mu_X \colon (X, \omega) \to B_X$ is a locally toric Lagrangian fibration, then B_X is equipped with a rigid structure called an *integral affine structure* (see Definition 7.1, also consult [11] and [32, Lemma 2.5] for more details). We show in Section 7 that there is a characteristic pair associated with an integral affine structure, and the canonical model constructed by this characteristic pair admits a symplectic structure so that the orbit map is a locally toric Lagrangian fibration. We shall refine the method used to prove Theorem 6.1 to obtain the classification theorem of locally toric Lagrangian fibrations (Theorem 7.8). This theorem has been obtained by Boucetta-Molino in [5].

This paper is organized as follows. In the next section we define the orbit space and the orbit map of a local torus action and investigate their properties. In Section 3 we see several examples. We give an obstruction in order that a local torus action is induced by a locally standard torus action. We also show that a locally toric Lagrangian fibration admits a C^∞ local torus action. In Section 4, we introduce a characteristic pair and construct a canonical model from a characteristic pair. We also explain how a characteristic pair is associated with a local torus action. In Section 5, we define the Euler class of the orbit map and prove that the orbit map has a C^0 section if and only if it vanishes. Section 6 is devoted to the topological classification. In Section 7 we investigate locally toric Lagrangian fibrations from the viewpoint of local torus actions. We give a necessary and sufficient condition in order that a manifold equipped with a local torus action admits a symplectic structure that makes the orbit map a locally toric Lagrangian fibration. We also reprove the classification theorem of locally toric Lagrangian fibrations by Boucetta-Molino. Finally, in Section 8, we investigate the topology such as fundamental groups, cohomology groups, and K-groups. In Appendix A we describe some facts about K-theory of low dimensional CW complexes which are used in this section. We also observe the signature in the oriented, four-dimensional case.

1.1. **Conventions.** Throughout this paper we employ the vector notation in order to represent elements of $\mathbb{C}^n$, namely, $z = (z_1, \dots, z_n) \in \mathbb{C}^n$. The similar notation is also used for $T^n = (S^1)^n$, $\mathbb{R}^n$, etc. In this paper, all manifolds, maps, and local T^n-actions are assumed to be of class C^0 unless otherwise stated.

Acknowledgment. I am thankful to Professor Margaret Symington for informing me of her work about Lagrangian fibrations and torus actions and to Professor Hisaaki Endo for explaining me Meyer's signature cocycle. I am also grateful to Professor Mikio Furuta and Professor Mikiya Masuda for helpful suggestions and for constant encouragement. Professor Furuta is my mentor and this work is motivated by his suggestion.

2. Orbit structures

2.1. **The standard representation.** In this subsection, we briefly summarize the fundamental facts about the standard representation of a torus, which is the local model in this paper. Recall that the standard representation of T^n is the T^n-action on $\mathbb{C}^n$ defined by

$$u \cdot z := (u_1 z_1, \dots, u_n z_n) \tag{2.1}$$

for $u = (u_1, \dots, u_n) \in T^n$ and $z = (z_1, \dots, z_n) \in \mathbb{C}^n$. The orbit space $\mathbb{C}^n/T^n$ of the action is equipped with the natural stratification whose k-dimensional strata consists of k-dimensional orbits. Let $\mathbb{R}^n_+$ be the standard n-dimensional positive cone which is defined by (1.3). It also has the natural stratification with respect to the number of components ξ_i of the coordinates $(\xi_1 \dots, \xi_n)$ which are equal to zero. We denote the k-dimensional part of $\mathbb{R}^n_+$ by $\mathcal{S}^{(k)}\mathbb{R}^n_+$. Namely, $\xi \in \mathbb{R}^n_+$ is an element of $\mathcal{S}^{(k)}\mathbb{R}^n_+$ if and only if the number of nonzero components ξ_i is equal to k. Then the map $\mu_{\mathbb{C}^n} \colon \mathbb{C}^n \to \mathbb{R}^n_+$ defined by (1.2) is invariant with respect to the standard representation of T^n and induces a homeomorphism from $\mathbb{C}^n/T^n$ to $\mathbb{R}^n_+$ which preserves stratifications. It is easy to see the following proposition.

Proposition 2.1. *For any $\xi \in \mathbb{R}^n_+$, all points of $\mu_{\mathbb{C}^n}^{-1}(\xi)$ has a common stabilizer. If ξ lies in $\mathcal{S}^{(k)}\mathbb{R}^n_+$, then it is an $(n-k)$-dimensional subtorus of T^n. In particular, for two points z_1, $z_2 \in \mathbb{C}^n$ with $\mu_{\mathbb{C}^n}(z_1) = \mu_{\mathbb{C}^n}(z_2)$, there exists an element $u \in T^n$ such that $z_2 = u \cdot z_1$. u is unique up to the common stabilizer of the T^n-action on $\mu_{\mathbb{C}^n}^{-1}(\mu_{\mathbb{C}^n}(z_1))$.*

Let ρ be an automorphism in $\mathrm{Aut}(T^n)$. Suppose that there exists a ρ-equivariant homeomorphism $f \colon \mathbb{C}^n \to \mathbb{C}^n$. Then, the following proposition is obvious.

Proposition 2.2. *For any $z \in \mathbb{C}^n$, ρ sends the stabilizer of z isomorphically to the one of $f(z)$.*

2.2. **orbit spaces and orbit maps.** Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^r local T^n-action. For $(X, \mathcal{T})$ we define the orbit space and the orbit map in the following way. Let $\{(U^X_\alpha, \varphi^X_\alpha)\}_{\alpha \in \mathcal{A}} \in \mathcal{T}$ be the maximal weakly standard atlas of X. We endow each quotient space $\varphi^X_\alpha(U^X_\alpha)/T^n$ with the quotient topology induced from the topology of $\varphi^X_\alpha(U^X_\alpha)$ by the natural projection $\pi_\alpha \colon \varphi^X_\alpha(U^X_\alpha) \to \varphi^X_\alpha(U^X_\alpha)/T^n$. By the property (ii) for each overlap $U^X_{\alpha\beta}$, $\varphi^X_{\alpha\beta}$ induces a homeomorphism from $\varphi^X_\beta(U^X_{\alpha\beta})/T^n$ to $\varphi^X_\alpha(U^X_{\alpha\beta})/T^n$. We define two elements $b_\alpha \in \varphi^X_\alpha(U^X_\alpha)/T^n$ and $b_\beta \in \varphi^X_\beta(U^X_\beta)/T^n$ to be equivalent, or $b_\alpha \sim_{orb} b_\beta$ if $b_\alpha \in \varphi^X_\alpha(U^X_{\alpha\beta})/T^n$, $b_\beta \in \varphi^X_\beta(U^X_{\alpha\beta})/T^n$ and the map induced by $\varphi^X_{\alpha\beta}$ sends b_β to b_α. It is an equivalence relation on the disjoint union $\coprod_\alpha \left(\varphi^X_\alpha(U^X_\alpha)/T^n\right)$. We define the *orbit space* B_X of the local T^n-action to be the quotient space

$$B_X := \coprod_{\alpha} \left(\varphi_\alpha^X(U_\alpha^X)/T^n\right) / \sim_{orb}$$

together with a quotient topology. It is easy to see that B_X is a Hausdorff space and $\{\varphi_\alpha^X(U_\alpha^X)/T^n\}_{\alpha\in\mathcal{A}}$ is an open covering of B_X. By the construction of B_X, the map $\coprod_\alpha \pi_\alpha \circ \varphi_\alpha^X \colon \coprod_\alpha U_\alpha^X \to \coprod_\alpha \left(\varphi_\alpha^X(U_\alpha^X)/T^n\right)$ induces a C^0 open map from X to B_X. We call it the *orbit map* of the local T^n-action and denote it by $\mu_X \colon X \to B_X$.

Definition 2.3 ([9, Section 6])**.** Let B be a Hausdorff space. A *structure of an n-dimensional C^0 manifold with corners* on B is a system of coordinate neighborhoods modeled on open subsets of $\mathbb{R}_+^n$ so that overlap maps are homeomorphisms which preserve the natural stratifications induced from the one of $\mathbb{R}_+^n$.

Proposition 2.4. *B_X is endowed with a structure of an n-dimensional C^0 manifold with corners.*

Proof. A structure of an n-dimensional C^0 manifold with corners on B_X is constructed as follows. We put $U_\alpha^B := \varphi_\alpha^X(U_\alpha^X)/T^n$. The restriction of $\mu_{\mathbb{C}^n}$ to $\varphi_\alpha^X(U_\alpha^X)$ induces a homeomorphism from U_α^B to the open subset $\mu_{\mathbb{C}^n}(\varphi_\alpha^X(U_\alpha^X))$ of $\mathbb{R}_+^n$, which is denoted by φ_α^B. By construction, on each overlap $U_{\alpha\beta}^B := U_\alpha^B \cap U_\beta^B$, the overlap map $\varphi_{\alpha\beta}^B := \varphi_\alpha^B \circ (\varphi_\beta^B)^{-1} \colon \mu_{\mathbb{C}^n}(\varphi_\beta^X(U_{\alpha\beta}^X)) \to \mu_{\mathbb{C}^n}(\varphi_\alpha^X(U_{\alpha\beta}^X))$ preserves the natural stratifications of $\mu_{\mathbb{C}^n}(\varphi_\alpha^X(U_{\alpha\beta}^X))$ and $\mu_{\mathbb{C}^n}(\varphi_\beta^X(U_{\alpha\beta}^X))$. Thus, $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ is the desired atlas. □

Remark 2.5. The atlas $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X constructed in the proof of Proposition 2.4 has the following property: for each α, $U_\alpha^X = \mu_X^{-1}(U_\alpha^B)$, $\varphi_\alpha^X(U_\alpha^X) = \mu_{\mathbb{C}^n}^{-1}(\varphi_\alpha^B(U_\alpha^B))$ and the following diagram commutes

$$\begin{tikzcd}
X \arrow[d, "\mu_X"] & \supset & \mu_X^{-1}(U_\alpha^B) \arrow[r, "\varphi_\alpha^X"] \arrow[d, "\mu_X"] & \mu_{\mathbb{C}^n}^{-1}(\varphi_\alpha^B(U_\alpha^B)) \arrow[d, "\mu_{\mathbb{C}^n}"] & \subset & \mathbb{C}^n \arrow[d, "\mu_{\mathbb{C}^n}"] \\
B_X & \supset & U_\alpha^B \arrow[r, "\varphi_\alpha^B"] & \varphi_\alpha^B(U_\alpha^B) & \subset & \mathbb{R}_+^n.
\end{tikzcd}$$

Remark 2.6. Let $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ be $2n$-dimensional manifolds equipped with C^r local T^n-actions. Suppose that there is a C^r isomorphism $f_X \colon (X_1, \mathcal{T}_1) \to (X_2, \mathcal{T}_2)$. Then, f_X induces a stratification preserving homeomorphism $f_B \colon B_{X_1} \to B_{X_2}$ such that f_X and f_B satisfy $\mu_{X_2} \circ f_X = f_B \circ \mu_{X_1}$. In Section 6, we classify local torus actions up to C^0 isomorphisms.

Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^r local T^n-action. By Proposition 2.4 B_X is equipped with a natural stratification. We denote by $\mathcal{S}^{(k)}B_X$ the k-dimensional part of B_X, namely, $\mathcal{S}^{(k)}B_X$ consists of those points which are sent to points in $\mathcal{S}^{(k)}\mathbb{R}_+^n$ by a local coordinate system. In particular, the top-dimensional part $\mathcal{S}^{(n)}B_X$ is equal to the interior of B_X.



Figure 1. The natural stratification when B_X is a triangle

Proposition 2.7. *For each k, the restriction $\mu_X|_{\mathcal{S}^{(k)}B_X}\colon \mu_X^{-1}(\mathcal{S}^{(k)}B_X)\to \mathcal{S}^{(k)}B_X$ of μ_X to $\mathcal{S}^{(k)}B_X$ admits a structure of a C^r fiber bundle whose fiber is T^k. In particular, $\mu_X|_{\mathcal{S}^{(n)}B_X}\colon \mu_X^{-1}(\mathcal{S}^{(n)}B_X)\to \mathcal{S}^{(n)}B_X$ is a C^r fiber bundle whose fiber is T^n and whose structure group is the semidirect product $T^n\rtimes \operatorname{Aut}(T^n)$ of T^n and $\operatorname{Aut}(T^n)$.*

Proof. It is sufficient to prove the proposition connected componentwise. Let $(\mathcal{S}^{(k)}B_X)_a$ ($a=1,\dots,m$) be the connected component of $\mathcal{S}^{(k)}B_X$. We take a weakly standard atlas $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}\in\mathcal{T}$. It induces an atlas $\{(U_\alpha^B,\varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X which satisfies the property in Remark 2.5. Let $(U_\alpha^B,\varphi_\alpha^B)$ be a coordinate neighborhood with $(\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B\neq\emptyset$. Then $(\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B$ is represented as common zeros of some exactly $n-k$ components, say $\varphi^B_{\alpha,i_1},\cdots,\varphi^B_{\alpha,i_{n-k}}$, of φ_α^B, namely,

$$(\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B=\{b\in U_\alpha^B\colon \varphi^B_{\alpha,i_1}(b)=\cdots=\varphi^B_{\alpha,i_{n-k}}(b)=0\}.$$

We put

$$T_{i_1,\dots,i_{n-k}}:=\{u\in T^n\colon u_j=1 \text{ unless } j\neq i_1,\dots,i_{n-k}\}.$$

Note that $T^n/T_{i_1,\dots,i_{n-k}}$ is identified with T^k. Let $\iota\colon\mathbb{R}^n_+\to\mathbb{C}^n$ be the section of $\mu_{\mathbb{C}^n}\colon\mathbb{C}^n\to\mathbb{R}^n_+$ which is defined by

$$\iota(\xi):=(\xi_1^{1/2},\dots,\xi_n^{1/2}). \tag{2.2}$$

By Proposition 2.1 the equation

$$\varphi_\alpha^X(x)=u_\alpha(x)\cdot\iota\circ\varphi_\alpha^B\circ\mu_X(x)$$

uniquely defines the map $u_\alpha\colon\mu_X^{-1}((\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B)\to T^n/T_{i_1,\dots,i_{n-k}}$. We define the local trivialization $\phi_\alpha\colon\mu_X^{-1}((\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B)\to(\mathcal{S}^{(k)}B_X)_a\cap U_\alpha^B\times T^k$ by

$$\phi_\alpha(x):=(\mu_X(x),u_\alpha(x)), \tag{2.3}$$

where we used the identification $T^n/T_{i_1,\dots i_{n-k}}\cong T^k$. This proves the first half. Next we prove the other part. For $k=n$, the direct computation shows that on a nonempty overlap $U^B_{\alpha\beta}\setminus\partial B_X$,

$$\phi_\alpha\circ\phi_\beta^{-1}(b,u)=(b,\rho_{\alpha\beta}(u)u_{\alpha\beta}(b))$$

for $(b,u)\in U^B_{\alpha\beta}\setminus\partial B_X\times T^n$, where $u_{\alpha\beta}$ is the map $u_{\alpha\beta}\colon U^B_{\alpha\beta}\setminus\partial B_X\to T^n$ which is defined uniquely by

$$\varphi^X_{\alpha\beta}\circ\iota\circ\varphi^B_\beta(b)=u_{\alpha\beta}(b)\cdot\iota\circ\varphi^B_\alpha(b). \tag{2.4}$$

This implies that the transition function with respect to ϕ_α and ϕ_β, which is denoted by $g_{\alpha\beta}$, is of the form $g_{\alpha\beta}(b)=(u_{\alpha\beta}(b),\rho_{\alpha\beta})\in T^n\rtimes\operatorname{Aut}(T^n)$. □

3. Examples

3.1. Locally standard torus actions.

Definition 3.1 (Locally standard torus actions)**.** Let X be a $2n$-dimensional C^∞ manifold equipped with a C^∞ T^n-action. A *standard coordinate neighborhood* of X is a triple (U,ρ,φ) consisting of a T^n-invariant open set U of X, an automorphism ρ of T^n, and a ρ-equivariant diffeomorphism φ from U to some T^n-invariant open subset in $\mathbb{C}^n$. The action of T^n on X is said to be *locally standard* if every point in X lies in some standard coordinate neighborhood and an atlas which consists of standard coordinate neighborhoods is called a *standard atlas.*

See [10, 6] for more details. A standard atlas is weakly standard. Therefore, a locally standard T^n-action on a closed C^∞ manifold X induces a local T^n-action on X. We give examples of locally standard torus actions.

Example 3.2. An effective C^∞ T^2-action on a 4-dimensional C^∞ manifold X without nontrivial finite stabilizers is locally standard because of the slice theorem. These actions have been studied by Orlik-Raymond in [31].

Example 3.3 (Nonsingular toric varieties)**.** A complex n-dimensional toric variety is a normal complex algebraic variety X of dimension n with a $(\mathbb{C}^*)^n$-action having a dense orbit. T^n acts on X as a subgroup of $(\mathbb{C}^*)^n$. If X is nonsingular, the T^n-action on X is locally standard. In fact, it is well-known that there is a one-to-one correspondence between toric varieties and fans, and top-dimensional cones in the fan associated with X correspond to standard coordinate neighborhoods all of which cover X since all cones are nonsingular. For toric varieties, see [8, 13, 30].

Example 3.4 (Quasi-toric manifolds)**.** A quasi-toric manifold is a C^∞ manifold equipped with a locally standard torus action whose orbit space is combinatorially isomorphic to a simple convex polytope. A quasi-toric manifold was first introduced by Davis-Januszkiewicz in their pioneer work [10] as a topological generalization of a projective toric variety and now, it plays a central role in toric topology. See [10, 6] for more details.

Note that an existence of a local T^n-action does not necessarily imply an existence of a locally standard T^n-action. For example, a nontrivial fiber bundle on an n-dimensional closed manifold whose fiber is T^n and whose structure group is $\mathrm{Aut}(T^n)$ is equipped with a local T^n-action which is not induced by any locally standard T^n-action. In general, for any C^r local T^n-action $\mathcal{T}$ on X, we take a weakly standard atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$. We denote by $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ the atlas of B_X which is induced by $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ and satisfies the property in Remark 2.5. Assume that the index set $\mathcal{A}$ is countable ordered. It is easy to see that the automorphisms $\rho_{\alpha\beta}$ of T^n in the property (ii) of Definition 1.1 form a Čech one-cocycle $\{\rho_{\alpha\beta}\}$ on $\mathcal{U} := \{U_\alpha^B\}_{\alpha\in\mathcal{A}}$ with values in $\mathrm{Aut}(T^n)$. By definition, two one-cocycles $\{\rho_{\alpha\beta}\}$ and $\{\rho'_{\alpha\beta}\}$ on $\mathcal{U}$ are equivalent if and only if there is a Čech zero-cochain $\{\rho_\alpha\}$ such that

$$\rho'_{\alpha\beta} = \rho_\alpha \circ \rho_{\alpha\beta} \circ \rho_\beta^{-1}$$

on each overlap $U_{\alpha\beta}^B$. We denote the set of equivalence classes by $H^1(\mathcal{U}; \mathrm{Aut}(T^n))$ and also denote its direct limit $\varinjlim H^1(\mathcal{U}; \mathrm{Aut}(T^n))$ with respect to refinements of $\mathcal{U}$ by $H^1(B_X; \mathrm{Aut}(T^n))$. See [24, Appendix A] for more details. $\{\rho_{\alpha\beta}\}$ determines an equivalence class in $H^1(B_X; \mathrm{Aut}(T^n))$. It does not depend on the choice of equivalent weakly standard atlases and depends only on the local T^n-action. The equivalence class of $\{\rho_{\beta\alpha}\}$ in $H^1(B_X; \mathrm{Aut}(T^n))$ is an obstruction for the local T^n-action to be induced by a locally standard T^n-action.

Proposition 3.5. *A C^r local T^n-action $\mathcal{T}$ on X is induced by some C^r locally standard T^n-action if and only if $\{\rho_{\alpha\beta}\}$ is equivalent to the trivial Čech one-cocycle in $H^1(B_X; \mathrm{Aut}(T^n))$, where the trivial Čech one-cocycle is the one whose values on all open set are equal to the identity map of T^n.*

Proof. If a C^r local T^n-action $\mathcal{T}$ on X is induced by some C^r locally standard T^n-action, then there exists a standard atlas $\{(U_\alpha^X, \rho_\alpha, \varphi_\alpha^X)\}$ in $\mathcal{T}$ such that $\rho_{\alpha\beta}$ is of the form $\rho_{\alpha\beta} = \rho_\alpha \circ \rho_\beta^{-1}$ on each overlap $U_{\alpha\beta}^B$. This implies that $\{\rho_{\alpha\beta}\}$ is equivalent to the trivial Čech one-cocycle.

Conversely, assume that $\{\rho_{\alpha\beta}\}$ is equivalent to the trivial Čech one-cocycle in $H^1(B_X; \mathrm{Aut}(T^n))$. By replacing the open covering of B_X by a refinement if necessary, we may assume that there exists a Čech zero-cochain $\{\rho_\alpha\}$ on $\{U_\alpha^B\}$ such that $\rho_\alpha \circ \rho_\beta^{-1} = \rho_{\alpha\beta}$ for each $U^B_{\alpha\beta}$. Then we can define the T^n-action on X by

$$u \cdot x := (\varphi^X_\alpha)^{-1}(\rho_\alpha(u) \cdot \varphi^X_\alpha(x))$$

for $u \in T^n$ and $x \in X$ provided that x lies in $\mu_X^{-1}(U^B_\alpha)$. The action is well-defined because if x also lies in $\mu_X^{-1}(U^B_\beta)$,

$$\begin{aligned}(\varphi^X_\alpha)^{-1}(\rho_\alpha(u) \cdot \varphi^X_\alpha(x)) &= (\varphi^X_\alpha)^{-1}(\rho_\alpha \circ \rho_\beta^{-1} \circ \rho_\beta(u) \cdot \varphi^X_\alpha \circ (\varphi^X_\beta)^{-1} \circ \varphi^X_\beta(x)) \\ &= (\varphi^X_\alpha)^{-1}(\rho_{\alpha\beta}(\rho_\beta(u)) \cdot \varphi^X_{\alpha\beta}(\varphi^X_\beta(x))) \\ &= (\varphi^X_\alpha)^{-1}(\varphi^X_{\alpha\beta}(\rho_\beta(u) \cdot \varphi^X_\beta(x))) \\ &= (\varphi^X_\beta)^{-1}(\rho_\beta(u) \cdot \varphi^X_\beta(x)),\end{aligned}$$

and for u_1, $u_2 \in T^n$,

$$\begin{aligned}u_1 \cdot (u_2 \cdot x) &= (\varphi^X_\alpha)^{-1}(\rho_\alpha(u_1) \cdot \varphi^X_\alpha((\varphi^X_\alpha)^{-1}(\rho_\alpha(u_2) \cdot \varphi^X_\alpha(x)))) \\ &= (\varphi^X_\alpha)^{-1}(\rho_\alpha(u_1u_2) \cdot \varphi^X_\alpha(x)) \\ &= (u_1u_2) \cdot x.\end{aligned}$$

□

We can construct an example of a local torus action which is not induced by any global torus action.

Example 3.6. Let B be a two-dimensional C^∞ torus with one boundary component and one corner points (see Figure 2). We construct a four-dimensional C^∞ manifold X equipped with a C^∞ local T^2-action whose orbit space is B as follows. First we focus on the interior $B \setminus \partial B$ of B which is denoted by B_1. Let $\rho\colon \pi_1(B)(\cong \pi_1(B_1)) \to SL_2(\mathbb{Z})$ be the representation of the fundamental group which is defined by



FIGURE 2. B

$$\rho([\alpha]) = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix},\ \rho([\beta]) = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix},\ \rho([\gamma]) = \begin{pmatrix} 3 & 1 \\ -1 & 0 \end{pmatrix},$$

where α, β, and γ are representatives of generators of $\pi_1(B)$ which satisfy the relation $[\alpha][\beta][\alpha]^{-1}[\beta]^{-1}[\gamma] = 1$ (see Figure 2). We identify T^2 with $\mathbb{R}^2/\mathbb{Z}^2$ and denote by $\pi_T\colon T^2_\rho \to B_1$ the associated T^2-bundle of the universal covering of B_1 by ρ.

Next we pay attention to a neighborhood of the boundary ∂B of B. We define subsets $\overline{B}_2$, U_1, and U_2 of $\mathbb{R}^2_+$ by

$$\begin{aligned}\overline{B}_2 &= \{\xi \in \mathbb{R}^2 \colon 0 \leq \xi_1 < 4,\ 0 \leq \xi_2 < 1\} \cup \{\xi \in \mathbb{R}^2 \colon 0 \leq \xi_1 < 1,\ 0 \leq \xi_2 < 4\},\\ U_1 &= \{\xi \in \mathbb{R}^2 \colon 3 < \xi_1 < 4,\ 0 \leq \xi_2 < 1\},\\ U_2 &= \{\xi \in \mathbb{R}^2 \colon 0 \leq \xi_1 < 1,\ 3 < \xi_2 < 4\},\end{aligned}$$

and also define diffeomorphisms $\varphi^B \colon U_1 \to U_2$ and $\varphi^X \colon \mu_{\mathbb{C}^2}^{-1}(U_1) \to \mu_{\mathbb{C}^2}^{-1}(U_2)$ by

$$\begin{aligned}\varphi^B(\xi) &= (\xi_2, 7 - \xi_1),\\ \varphi^X(z) &= \left(\frac{z_1^3 z_2}{|z_1|^3}, \sqrt{7 - |z_1|^2}\left(\frac{z_1}{|z_1|}\right)^{-1}\right).\end{aligned}$$

Note that φ^B and φ^X satisfy

$$\mu_{\mathbb{C}^2} \circ \varphi^X = \varphi^B \circ \mu_{\mathbb{C}^2}. \tag{3.1}$$

We denote by X_2 the manifold which is obtained from $\mu_{\mathbb{C}^2}^{-1}(\overline{B}_2)$ by gluing $\mu_{\mathbb{C}^2}^{-1}(U_1)$ and $\mu_{\mathbb{C}^2}^{-1}(U_2)$ with φ^X and denote by B_2 the surface with one corner which is obtained from $\overline{B}_2$ by gluing U_1 and U_2 with φ^B. B_2 can be identified with a neighborhood of ∂B. By (3.1), $\mu_{\mathbb{C}^2}$ descends to the map from X_2 to B_2. We denote it by $\mu_2 \colon X_2 \to B_2$. It is easy to see that the restriction $\mu_2|_{\mu_2^{-1}(B_1 \cap B_2)} \colon \mu_2^{-1}(B_1 \cap B_2) \to B_1 \cap B_2$ is a fiber bundle with fiber T^2 and structure group $SL_2(\mathbb{Z})$ which is isomorphic to $\pi_T|_{B_1 \cap B_2} \colon T^2_\rho|_{B_1 \cap B_2} \to B_1 \cap B_2$ because they have the same monodromy. Thus we can glue $\pi_T \colon T^2_\rho \to B_1$ and $\mu_2 \colon X_2 \to B_2$ together to get the map $\mu \colon X \to B$. By the construction, X is a C^∞ four-dimensional manifold equipped with a local T^2-action whose orbit space B_X and orbit map μ_X are equal to B and μ, respectively. Note that μ is C^∞ in this example.



Figure 3. B_2

3.2. **Locally toric Lagrangian fibrations.**

Let (X, ω) be a $2n$-dimensional closed C^∞ symplectic manifold and B an n-dimensional C^∞ manifold with corners.

Definition 3.7 (Locally toric Lagrangian fibrations [19, Definition 2.7])**.** A C^∞ map $\mu \colon (X, \omega) \to B$ is called a *locally toric Lagrangian fibration* if there exists a system $\{(U_\alpha, \varphi^B_\alpha)\}$ of coordinate neighborhoods of B modeled on $\mathbb{R}^n_+$, and for each α there exists a symplectomorphism $\varphi^X_\alpha \colon (\mu^{-1}(U_\alpha), \omega) \to (\mu_{\mathbb{C}^n}^{-1}(\varphi^B_\alpha(U_\alpha)), \omega_{\mathbb{C}^n})$ such that $\mu_{\mathbb{C}^n} \circ \varphi^X_\alpha = \varphi^B_\alpha \circ \mu$.

Example 3.8. A moment map of a symplectic toric manifold is a locally toric Lagrangian fibration.

Example 3.9. Let $\omega_{\mathbb{R}^n \times T^n}$ be the standard symplectic structure on $\mathbb{R}^n \times T^n$ which is defined by

$$\omega_{\mathbb{R}^n \times T^n} = \sum_{k=1}^n d\theta_k \wedge d\xi_k, \tag{3.2}$$

where $(\xi_1, \dots, \xi_n)$ is the standard coordinates of $\mathbb{R}^n$ and $(\theta_1, \dots, \theta_n)$ is the angle coordinates of T^n with period 1, which means $(e^{2\pi\theta_1}, \dots, e^{2\pi\theta_n}) \in T^n$. Then the projection $\mathrm{pr}_1 \colon (\mathbb{R}^n \times T^n, \omega_{\mathbb{R}^n \times T^n}) \to \mathbb{R}^n$ to the first factor is a nonsingular Lagrangian fibration. It is easy to see that $\mathrm{pr}_1 \colon (\mathbb{R}^n \times T^n, \omega_{\mathbb{R}^n \times T^n}) \to \mathbb{R}^n$ is a noncompact locally toric Lagrangian fibration.

The Arnold-Liouville theorem [1] says that $\mathrm{pr}_1 \colon (\mathbb{R}^n \times T^n, \omega_{\mathbb{R}^n \times T^n}) \to \mathbb{R}^n$ is the local model of a nonsingular Lagrangian fibration.

Theorem 3.10 (Arnold-Liouville [1])**.** *Let $\mu \colon (X, \omega) \to B$ be a nonsingular Lagrangian fibration with closed connected fibers on a closed manifold. Then for each $b \in B$ there exists a coordinate neighborhood (U, φ^B) and there also exists a symplectomorphism $\varphi^X \colon (\mu^{-1}(U), \omega) \to (\varphi^B(U) \times T^n, \omega_{\mathbb{R}^n \times T^n})$ such that $\mathrm{pr}_1 \circ \varphi^X = \varphi^B \circ \mu$.*

Therefore, a nonsingular Lagrangian fibration with closed connected fibers on a closed manifold is also a locally toric Lagrangian fibration. Nonsingular Lagrangian fibrations on closed oriented surfaces have been investigated by Mishachev. In [29] he showed that the complete list of nonsingular Lagrangian fibrations on closed oriented surfaces. According to it, if a closed oriented surface B is a base of a nonsingular Lagrangian fibration, then B is T^2. Moreover, concrete constructions of all these have been given in [16].

Example 3.11. We can construct an example of locally toric Lagrangian fibrations as follows. For a sufficiently small positive number ε $(1 \gg \varepsilon > 0)$, let U be the open subset of $(\mathbb{C}^2, \omega_{\mathbb{C}^2})$ which is defined by

$$U := \{z \in \mathbb{C}^2 \colon z_1 \neq 0, |z_2|^2 > |z_1|^2 - (1+\varepsilon)\}.$$

S^1 acts on $(U, \omega_{\mathbb{C}^2})$ by

$$u \cdot z := (z_1, u^{-1} z_2)$$

for $u \in S^1$ and $z \in U$. This action is Hamiltonian and the map $\mu_{S^1} \colon U \to \mathbb{R}$ defined by

$$\mu_{S^1}(z) := 1 - |z_2|^2$$

is a moment map of this action. We denote by $\overline{X}$ the cut space by the symplectic cutting with respect to the S^1-action, namely, $\overline{X}$ is the quotient space

$$\overline{X} := \{(z, w) \in U \times \mathbb{C} \colon 1 - |z_2|^2 - |w|^2 = 0\}/S^1$$

by the S^1-action

$$u \cdot (z, w) := (z_1, u^{-1} z_2, u^{-1} w).$$

By [25], $\overline{X}$ is a symplectic manifold and the restriction of the standard T^2-action on $\mathbb{C}^2$ to U induces a Hamiltonian T^2-action on $\overline{X}$ with a moment map $\overline{\mu}([z, w]) := (|z_1|^2, |z_2|^2)$. We denote by $\overline{B}$ the image of $\overline{\mu}$. $\overline{B}$ is written by

$$\overline{B} = \{\xi \in \mathbb{R}^2 \colon \xi_1 > 0, 0 \leq \xi_2 \leq 1, \xi_2 > \xi_1 - (1+\varepsilon)\}.$$

Let $\overline{X}_1$ and $\overline{X}_2$ be the open subspace of $\overline{X}$ which are defined by

$$\overline{X}_1 := \{[z, w] \in \overline{X} \colon |z_1|^2 < \varepsilon\},$$
$$\overline{X}_2 := \{[z, w] \in \overline{X} \colon |z_1|^2 - (1+\varepsilon) < |z_2|^2 < |z_1|^2 - 1\}.$$

We define the diffeomorphism $\varphi^X \colon \overline{X}_1 \to \overline{X}_2$ by

$$\varphi^X([z, w]) := \left[\left((z_1/|z_1|)\sqrt{|z_1|^2 + |z_2|^2 + 1}, (z_1/|z_1|)^{-1} z_2\right), w\right].$$

It is easy to see that φ^X is well-defined and preserves symplectic structures of $\overline{X}_1$ and $\overline{X}_2$ induced by the symplectic structure on $\overline{X}$. The images of $\overline{X}_1$ and $\overline{X}_2$ by $\overline{\mu}$ are open subset of $\overline{B}$ which are written by

$$\overline{\mu}(\overline{X}_1) = \{\xi \in \overline{B} \colon \xi_1 < \varepsilon\}, \quad \overline{\mu}(\overline{X}_2) = \{\xi \in \overline{B} \colon \xi_2 < \xi_1 - 1\}.$$

φ^X covers the diffeomorphism $\varphi^B\colon \overline{\mu}(\overline{X}_1) \to \overline{\mu}(\overline{X}_2)$ by

$$\varphi^B(\xi) := (\xi_1 + \xi_2 + 1, \xi_2).$$

This means that they satisfy

$$\overline{\mu} \circ \varphi^X = \varphi^B \circ \overline{\mu}. \tag{3.3}$$

We define X to be the quotient space

$$X := \overline{X}/\sim_X,$$

where $[z_1, w_1] \sim_X [z_2, w_2]$ if and only if $[z_i, w_i] \in \overline{X}_i$ for $i = 1, 2$ and $[z_2, w_2] = \varphi^X([z_1, w_1])$ and also define B to be the quotient space

$$B := \overline{B}/\sim_B,$$

where $\xi \sim_B \eta$ if and only if $\xi \in \overline{\mu}(\overline{X}_1)$, $\eta \in \overline{\mu}(\overline{X}_2)$, and $\eta = \varphi^B(\xi)$. Since $\overline{\mu}$ satisfies (3.3), it descends to the map $\mu\colon X \to B$. By the construction, the symplectic structure on $\overline{X}$ induces a symplectic structure ω on X so that $\mu\colon (X, \omega) \to B$ is a locally toric Lagrangian fibration.

We show that a locally toric Lagrangian fibration is endowed with a C^∞ local torus action. First we recall the following fact about automorphisms of the local models of a nonsingular Lagrangian fibration. In this lemma we identify an automorphism of T^n with an element of $\mathrm{GL}_n(\mathbb{Z})$.

Lemma 3.12 ([32, Lemma 2.5]). *Let φ be a fiber-preserving symplectomorphism of $\mathrm{pr}_1\colon (\mathbb{R}^n \times T^n, \omega_{\mathbb{R}^n\times T^n}) \to \mathbb{R}^n$. Then, φ is of the form $\varphi(\xi, u) = (\rho^{-T}(\xi) + c, \rho(u)f(\xi))$ for some $\rho \in \mathrm{Aut}(T^n)$, $c \in \mathbb{R}^n$ and a Lagrangian section f of $\mathrm{pr}_1\colon (\mathbb{R}^n\times T^n, \omega_{\mathbb{R}^n\times T^n}) \to \mathbb{R}^n$, where ρ^{-T} is the transpose inverse of ρ. The latter means that $f^*\omega_{\mathbb{R}^n\times T^n}$ vanishes.*

We show the following proposition.

Proposition 3.13. *Let $\mu\colon (X, \omega) \to B$ be a locally toric Lagrangian fibration on an n-dimensional base B and $\{(U_\alpha, \varphi^B_\alpha, \varphi^X_\alpha)\}$ a system of local identifications of μ with $\mu_{\mathbb{C}^n}$. Then, on each nonempty overlap $U_{\alpha\beta} := U_\alpha \cap U_\beta$, there exists an element $\rho_{\alpha\beta} \in \mathrm{Aut}(T^n)$ such that $\varphi^X_\alpha \circ (\varphi^X_\beta)^{-1}$ is $\rho_{\alpha\beta}$-equivariant.*

Proof. First we focus on the interior $B\backslash\partial B$ of B. Since the restriction of $\mu\colon (X, \omega) \to B$ to $B\backslash\partial B$ is a nonsingular Lagrangian fibration, it is locally identified with the local model. In fact, we can construct local identifications explicitly. For each α, φ^X_α sends $(\mu^{-1}(U_\alpha \setminus \partial B), \omega)$ fiber-preserving symplectomorphically to $(\mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha \setminus \partial B)), \omega_{\mathbb{C}^n})$ which covers φ^B_α. We define a fiber preserving symplectomorphism $\nu_{\mathbb{C}^n}\colon (\varphi^B_\alpha(U_\alpha \setminus \partial B) \times T^n, \omega_{\mathbb{R}^n\times T^n}) \to (\mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_\alpha \setminus \partial B)), \omega_{\mathbb{C}^n})$ by

$$\nu_{\mathbb{C}^n}(\xi, u) := u \cdot \iota(\xi), \tag{3.4}$$

where ι is the map defined by (2.2). Note that T^n acts on $\varphi^B_\alpha(U_\alpha \setminus \partial B) \times T^n$ by the multiplication to the second factor and $\nu_{\mathbb{C}^n}$ is equivariant with respect to this action and the standard representation of T^n. Thus $\nu^{-1}_{\mathbb{C}^n} \circ \varphi^X_\alpha$ is a local identification.

On each nonempty overlap $U_{\alpha\beta}$, by applying Lemma 3.12 to the fiber preserving symplectomorphism $\nu^{-1}_{\mathbb{C}^n} \circ \varphi^X_\alpha \circ (\varphi^X_\beta)^{-1} \circ \nu_{\mathbb{C}^n}\colon (\varphi^B_\beta(U_{\alpha\beta}\setminus\partial B), \omega_{\mathbb{R}^n\times T^n}) \to (\varphi^B_\alpha(U_{\alpha\beta}\setminus \partial B), \omega_{\mathbb{R}^n\times T^n})$, on $U_{\alpha\beta}$ there exists an automorphism $\rho_{\alpha\beta} \in \mathrm{Aut}(T^n)$ such that $\nu^{-1}_{\mathbb{C}^n} \circ \varphi^X_\alpha \circ (\varphi^X_\beta)^{-1} \circ \nu_{\mathbb{C}^n}$ is $\rho_{\alpha\beta}$-equivariant. Since $\nu_{\mathbb{C}^n}$ is equivariant and $\mu^{-1}(U_{\alpha\beta} \setminus \partial B)$ is open dense in $\mu^{-1}(U_{\alpha\beta})$, the overlap map $\varphi^X_\alpha \circ (\varphi^X_\beta)^{-1}\colon \mu^{-1}_{\mathbb{C}^n}(\varphi^B_\beta(U_{\alpha\beta})) \to \mu^{-1}_{\mathbb{C}^n}(\varphi^B_\alpha(U_{\alpha\beta}))$ is also $\rho_{\alpha\beta}$-equivariant. □

This implies that $\{(\mu^{-1}(U_\alpha), \varphi_\alpha^X)\}$ is a weakly standard atlas of X. Hence, X is equipped with a C^∞ local T^n-action. It is obvious that the orbit space and the orbit map of the local T^n-action on X are naturally identified with B and μ. In Section 7, we will see a necessary and sufficient condition in order that a manifold with a local torus action becomes a locally toric Lagrangian fibration.

3.3. **New local torus actions from given ones.** When a local torus action is given, we can construct new local torus actions from the given one.

Example 3.14 (fiber products)**.** Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^r local T^n-action. Suppose that $f\colon B' \to B_X$ is a stratification preserving locally homeomorphism from an n-dimensional C^0 manifold B' with corners to B_X. Then it is obvious that the fiber product $f^*X := \{(b', x) \in B' \times X \colon f(b') = \mu_X(x)\}$ of f and μ_X is equipped with a C^r local T^n-action whose orbit space is B'.

Example 3.15 (blowing-ups)**.** Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^∞ local T^n-action. Let $x \in X$. Suppose that there exists a coordinate neighborhood (U^X, φ^X) in a weakly standard atlas of $\mathcal{T}$ such that φ^X sends x to the origin of $\mathbb{C}^n$. For a sufficiently small positive real number $\varepsilon > 0$, we denote by $\varphi^X(U^X)_{\geq\varepsilon}$ the quotient space of $\{(z, w) \in \varphi^X(U^X) \times \mathbb{C} \colon \|z\|^2 - |w|^2 = \varepsilon\}$ by the circle action defined by

$$t \cdot (z, w) := (tz_1, \ldots, tz_n, t^{-1}w).$$

It is easy to see that $\varphi^X(U^X)_{\geq\varepsilon}$ is smooth and T^n acts on $\varphi^X(U^X)_{\geq\varepsilon}$ by

$$u \cdot [z, w] := [u \cdot z, w].$$

Let $\overline{D}_\varepsilon(0)$ be the closed disc in $\mathbb{C}^n$ centered at the origin with radius $\varepsilon^{1/2}$. We define the diffeomorphism f_ε from the open set $\varphi^X(U^X) \setminus \overline{D}_\varepsilon(0)$ to $\{[z, w] \in \varphi^X(U^X)_{\geq\varepsilon} \colon w \neq 0\}$ by

$$f_\varepsilon(z) = \left[z, (\|z\|^2 - \varepsilon)^{1/2}\right]$$

for $z \in \varphi^X(U^X) \setminus \overline{D}_\varepsilon(0)$. It is easy to see that f_ε is an equivariantly diffeomorphism. By removing $(\varphi_\alpha^X)^{-1}(\overline{D}_\varepsilon(0))$ for a sufficiently small $\varepsilon > 0$ from X and gluing $X \setminus (\varphi_\alpha^X)^{-1}(\overline{D}_\varepsilon(0))$ and $\varphi^X(U^X)_{\geq\varepsilon}$ by $f_\varepsilon \circ \varphi_\alpha^X$, we can obtain a new manifold equipped with a C^∞ local T^n-action. We call it the *blowing-up* of X at x. This is a C^∞ aspect of the symplectic blowing-up by Guillemin-Sternberg in [17].

Example 3.16 (connected sums)**.** Let $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ be $2n$-dimensional manifolds equipped with C^r local T^n-actions. Suppose that (U^{X_1}, φ^{X_1}) and (U^{X_2}, φ^{X_2}) are coordinate neighborhoods in weakly standard atlases of $\mathcal{T}_1$ and $\mathcal{T}_2$ such that both φ^{X_1} and φ^{X_2} send U^{X_1} and U^{X_2} C^r diffeomorphically to a same T^n-invariant open set of $\mathbb{C}^n$. Then we can perform the connected sum $X_1 \# X_2$ and the obtained manifold $X_1 \# X_2$ has a C^r local T^n-action whose orbit space is the connected sum $B_{X_1} \# B_{X_2}$.

4. Characteristic pairs and canonical models

In this section, we introduce a characteristic pair and construct a canonical model from a characteristic pair. Both of them will play important roles in the rest of this paper. We also show that a characteristic pair is associated with a local torus action.

4.1. **Characteristic pairs.** Let B be an n-dimensional C^0 manifold with corners. We assume that $\partial B \neq \emptyset$. By the definition of a manifold with corners, B is equipped with a natural stratification. We denote by $\mathcal{S}^{(k)}B$ the k-dimensional part of B. Let Λ be the lattice of integral elements in the Lie algebra $\mathfrak{t}$ of T^n, namely,

$$\Lambda := \{t \in \mathfrak{t}\colon\ \exp t = 1\}.$$

Since the differential of any automorphism of T^n at the unit element preserves Λ, by assigning to an automorphism of T^n its differential at the unit element, there is a natural homomorphism from $\mathrm{Aut}(T^n)$ to $\mathrm{GL}(\Lambda)$. It is an isomorphism. In fact, it follows from the surjectivity of the exponential map of T^n and the equation $\varphi \circ \exp = \exp \circ d\varphi$ for any automorphism $\varphi \in \mathrm{Aut}(T^n)$. In the rest of this paper, we identify $\mathrm{Aut}(T^n)$ with $\mathrm{GL}(\Lambda)$ by this isomorphism. Let $\pi_P\colon P \to B$ be a principal $\mathrm{Aut}(T^n)$-bundle on B and $\pi_\Lambda\colon \Lambda_P \to B$ the associated Λ-bundle of P by the above isomorphism $\mathrm{Aut}(T^n) \cong \mathrm{GL}(\Lambda)$. Suppose that $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ is a rank one subbundle of the restriction $\pi_\Lambda|_{\mathcal{S}^{(n-1)}B}\colon \Lambda_P|_{\mathcal{S}^{(n-1)}B} \to \mathcal{S}^{(n-1)}B$ of $\pi_\Lambda\colon \Lambda_P \to B$ to $\mathcal{S}^{(n-1)}B$. For each k and any point $b \in \mathcal{S}^{(k)}B$, let U be an open neighborhood of b in B on which there exists a local trivialization $\varphi^\Lambda\colon \pi_\Lambda^{-1}(U) \to U \times \Lambda$ of Λ_P. By shrinking U if necessary, we can assume that the intersection $U \cap \mathcal{S}^{(n-1)}B$ has exactly $n-k$ connected components, say, $(U \cap \mathcal{S}^{(n-1)}B)_1, \ldots, (U \cap \mathcal{S}^{(n-1)}B)_{n-k}$. Since Λ is discrete, for each $(U \cap \mathcal{S}^{(n-1)}B)_a$ there exists a rank one sublattice $L_a \subset \Lambda$ such that φ^Λ sends $\pi_{\mathcal{L}}^{-1}\left((U \cap \mathcal{S}^{(n-1)}B)_a\right)$ fiberwise to $(U \cap \mathcal{S}^{(n-1)}B)_a \times L_a$


U
$(U \cap \mathcal{S}^{(1)}B)_1$
b
$(U \cap \mathcal{S}^{(1)}B)_2$

FIGURE 4. An open neighborhood U when B is a triangle and $b \in \mathcal{S}^{(0)}B$

$$\begin{array}{ccc}
\pi_\Lambda^{-1}(U) & \overset{\varphi^\Lambda}{\cong} & U \times \Lambda \\
\cup & & \cup \\
\pi_\Lambda^{-1}\left((U \cap \mathcal{S}^{(n-1)}B)_a\right) & \cong & (U \cap \mathcal{S}^{(n-1)}B)_a \times \Lambda \\
\cup & & \cup \\
\pi_{\mathcal{L}}^{-1}\left((U \cap \mathcal{S}^{(n-1)}B)_a\right) & \cong & (U \cap \mathcal{S}^{(n-1)}B)_a \times L_a.
\end{array}$$

Definition 4.1. $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ is said to be *unimodular* if for each k and any point $b \in \mathcal{S}^{(k)}B$, the sublattice $L_1 + \cdots + L_{n-k}$ generated by the above rank one sublattices $L_1, \ldots, L_{n-k}$ is a rank $n-k$ direct summand of Λ. (In [10] such a sublattice is called an $(n-k)$-*dimensional unimodular subspace* of Λ.)

Of course, $L_1, \ldots, L_{n-k}$ themselves depend on the choice of a neighborhood U and a local trivialization φ^Λ, but Definition 4.1 does not depend on the choice of them because unimodularness is invariant by an automorphism of Λ.

Definition 4.2. Let $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ be a unimodular rank one subbundle of $\pi_\Lambda|_{\mathcal{S}^{(n-1)}B}\colon \Lambda_P|_{\mathcal{S}^{(n-1)}B} \to \mathcal{S}^{(n-1)}B$. Then the pair $(P, \mathcal{L})$ of $\pi_P\colon P \to B$ and $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ is called a *characteristic pair* and $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ is called a *characteristic bundle.*

Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a local T^n-action. We show that there is a characteristic pair associated with $(X, \mathcal{T})$. Let $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ be the maximal weakly standard atlas. It induces an atlas $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X which satisfies the property in Remark 2.5 and also determines a Čech one-cocycle $\{\rho_{\alpha\beta}\}$ on $\{U_\alpha^B\}_{\alpha\in\mathcal{A}}$ with coefficients in $\operatorname{Aut}(T^n)$. It defines the principal $\operatorname{Aut}(T^n)$-bundle $\pi_{P_X} : P_X \to B_X$ on B_X by setting

$$P_X := \left(\coprod_\alpha U_\alpha^B \times \operatorname{Aut}(T^n)\right) / \sim_P, \tag{4.1}$$

where $(b_\alpha, h_\alpha) \in U_\alpha^B \times \operatorname{Aut}(T^n) \sim_P (b_\beta, h_\beta) \in U_\beta^B \times \operatorname{Aut}(T^n)$ if and only if $b_\alpha = b_\beta \in U_{\alpha\beta}^B$ and $h_\alpha = \rho_{\alpha\beta} \circ h_\beta$. The bundle projection π_{P_X} is defined by the obvious way. For each α, every point in $\pi_{P_X}^{-1}(U_\alpha^B)$ has a unique representative which lies in $U_\alpha^B \times \operatorname{Aut}(T^n)$. By associating a point in $\pi_{P_X}^{-1}(U_\alpha^B)$ with the unique representative, we define the local trivialization of P_X on U_α^B which is denoted by $\varphi_\alpha^P : \pi_{P_X}^{-1}(U_\alpha^B) \to U_\alpha^B \times \operatorname{Aut}(T^n)$. Note that the following equality holds for $(b, h) \in U_{\alpha\beta}^B \times \operatorname{Aut}(T^n)$

$$\varphi_{\alpha\beta}^P(b, h) := \varphi_\alpha^P \circ (\varphi_\beta^P)^{-1}(b, h) = (b, \rho_{\alpha\beta} \circ h). \tag{4.2}$$

We denote by $\pi_{\Lambda_X} : \Lambda_X \to B_X$ the Λ-bundle associated with P_X. We show that the property (ii) in Definition 1.1 determines a unique unimodular rank one subbundle of $\pi_{\Lambda_X}|_{\mathcal{S}^{(n-1)}B_X} : \Lambda_X|_{\mathcal{S}^{(n-1)}B_X} \to \mathcal{S}^{(n-1)}B_X$. For each coordinate neighborhood $(U_\alpha^B, \varphi_\alpha^B)$ of B_X with $U_\alpha^B \cap \mathcal{S}^{(n-1)}B_X \neq \emptyset$, T^n acts on the preimage $\mu_{\mathbb{C}^n}^{-1}\left(\varphi_\alpha^B(U_\alpha^B \cap \mathcal{S}^{(n-1)}B_X)\right)$ as the restriction of the standard representation of T^n. For simplicity, we assume that the intersection $U_\alpha^B \cap \mathcal{S}^{(n-1)}B_X$ is connected. (Otherwise, we may consider componentwise.) Then, $\mu_{\mathbb{C}^n}^{-1}\left(\varphi_\alpha^B(U_\alpha^B \cap \mathcal{S}^{(n-1)}B_X)\right)$ is fixed by a circle subgroup of T^n. We denote the circle subgroup by S_α^1 and also denote the rank one sublattice of Λ spanned by the integral element which generates S_α^1 by $\mathcal{L}_\alpha$. Suppose that $(U_\alpha^B, \varphi_\alpha^B)$ and $(U_\beta^B, \varphi_\beta^B)$ are coordinate neighborhoods satisfying the above conditions and the intersection $U_{\alpha\beta}^B \cap \mathcal{S}^{(n-1)}B_X$ is nonempty.

Lemma 4.3. *Under the identification* $\operatorname{Aut}(T^n) \cong \operatorname{GL}(\Lambda)$*, the automorphism* $\rho_{\alpha\beta}$ *of* T^n *in the property* (ii) *of Definition* 1.1 *sends* $\mathcal{L}_\beta$ *isomorphically to* $\mathcal{L}_\alpha$*.*

Proof. Since $\varphi_{\alpha\beta}^X$ is a $\rho_{\alpha\beta}$-equivariant homeomorphism, the automorphism $\rho_{\alpha\beta}$ of T^n sends S_β^1 isomorphically to S_α^1. See Proposition 2.2. Therefore, $\rho_{\alpha\beta}$ also sends $\mathcal{L}_\beta$ isomorphically to $\mathcal{L}_\alpha$. □

By the construction of $\pi_{\Lambda_X} : \Lambda_X \to B_X$, there exists a local trivialization $\varphi_\alpha^\Lambda : \pi_{\Lambda_X}^{-1}(U_\alpha^B) \to U_\alpha^B \times \Lambda$ of $\pi_{\Lambda_X} : \Lambda_X \to B_X$ on each U_α^B such that on an overlap $U_{\alpha\beta}^B$ the transition function with respect to φ_α^Λ and φ_β^Λ is $\rho_{\alpha\beta}$. We take a subsystem $\{(U_{\alpha_i}^B, \varphi_{\alpha_i}^B)\}_{i\in\mathcal{I}}$ of $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ which covers $\mathcal{S}^{(n-1)}B_X$. By Lemma 4.3, we can obtain the rank one subbundle $\pi_{\mathcal{L}_X} : \mathcal{L}_X \to \mathcal{S}^{(n-1)}B_X$ of $\pi_{\Lambda_X}|_{\mathcal{S}^{(n-1)}B_X} : \Lambda_X|_{\mathcal{S}^{(n-1)}B_X} \to \mathcal{S}^{(n-1)}B_X$ by setting

$$\mathcal{L}_X := \left(\coprod_i U_{\alpha_i}^B \cap \mathcal{S}^{(n-1)}B_X \times \mathcal{L}_{\alpha_i}\right) / \sim_L, \tag{4.3}$$

where $(b_i, l_i) \in U_{\alpha_i}^B \cap \mathcal{S}^{(n-1)}B_X \times \mathcal{L}_{\alpha_i} \sim_L (b_j, l_j) \in U_{\alpha_j}^B \cap \mathcal{S}^{(n-1)}B_X \times \mathcal{L}_{\alpha_j}$ if and only if $b_i = b_j$ and $l_i = \rho_{\alpha_i\alpha_j}(l_j)$. By the construction, it is easy to see that $\pi_{\mathcal{L}_X} : \mathcal{L}_X \to \mathcal{S}^{(n-1)}B_X$ is unimodular. As a summary, we have the following proposition.

Proposition 4.4. *Associated with a local T^n-action $\mathcal{T}$ on X, there exists a characteristic pair $(P_X, \mathcal{L}_X)$, where P_X and $\mathcal{L}_X$ are defined by* (4.1) *and* (4.3)*, respectively.*

Example 4.5. For a $2n$-dimensional manifold X equipped with a locally standard T^n-action, $\pi_{P_X} \colon P_X \to B_X$ is the product principal $\mathrm{Aut}(T^n)$-bundle $P_X = B_X \times \mathrm{Aut}(T^n)$. Let $(\mathcal{S}^{(n-1)}B_X)_a$ $(a = 1, \dots, k)$ be the connected component of $\mathcal{S}^{(n-1)}B_X$. Then each preimage $\mu_X^{-1}((\mathcal{S}^{(n-1)}B_X)_a)$ is fixed by a circle subgroup S^1_a of T^n. Let L_a be the rank one sublattice in Λ corresponding to S^1_a. Then, $\mathcal{L}_X$ is the disjoint union $\coprod_a (\mathcal{S}^{(n-1)}B_X)_a \times L_a$. In particular, if X is a quasi-toric manifold, $\mathcal{L}_X$ is essentially nothing but the characteristic function. Since $\mathcal{L}_X$ is unimodular for a locally standard T^n-action on X, it is easy to see that B_X is *nice*, namely, each connected component of a k-dimensional part $\mathcal{S}^{(k)}B$ of B is contained in exactly $n-k$ facets.

Example 4.6. In the case of Example 3.6, we identify Λ with $\mathbb{Z} \oplus \mathbb{Z}$ and also identify $\mathrm{Aut}(T^2)$ with $\mathrm{GL}_2(\mathbb{Z})$. Then, $\pi_{P_X} \colon P_X \to B$ is the principal $\mathrm{Aut}(T^2)$-bundle associated with the universal covering of B by ρ. The restriction of $\pi_{\Lambda_X} \colon \Lambda_X \to B$ to $\mathcal{S}^{(1)}B$ (in this case, $n = 2$) and the characteristic bundle $\pi_{\mathcal{L}_X} \colon \mathcal{L}_X \to \mathcal{S}^{(1)}B$ are obtained by gluing two pairs

$$\left(\{\xi \in \overline{B}_2 \colon 0 < \xi_1,\ \xi_2 = 0\} \times \Lambda, \{\xi \in \overline{B}_2 \colon 0 < \xi_1,\ \xi_2 = 0\} \times (\{0\} \oplus \mathbb{Z})\right)$$

and

$$\left(\{\xi \in \overline{B}_2 \colon \xi_1 = 0,\ 0 < \xi_2\} \times \Lambda, \{\xi \in \overline{B}_2 \colon \xi_1 = 0,\ 0 < \xi_2\} \times (\mathbb{Z} \oplus \{0\})\right)$$

of trivial Λ-bundles and their rank one subbundles with the map $\varphi_\Lambda \colon \{\xi \in U_1 \colon 0 < \xi_1,\ \xi_2 = 0\} \times \Lambda \to \{\xi \in U_2 \colon \xi_1 = 0,\ 0 < \xi_2\} \times \Lambda$ defined by $\varphi_\Lambda(\xi, \lambda) := (\varphi_B(\xi), \rho(\gamma)(\lambda))$.

For $i = 1, 2$, let B_i be an n-dimensional C^0 manifold with corners and $(P_i, \mathcal{L}_i)$ a characteristic pair on B_i.

Definition 4.7. An *isomorphism* $f_P \colon (P_1, \mathcal{L}_1) \to (P_2, \mathcal{L}_2)$ between characteristic pairs is a bundle isomorphism $f_P \colon P_1 \to P_2$ which covers a stratification preserving homeomorphism $f_B \colon B_1 \to B_2$ such that the lattice bundle isomorphism $f_\Lambda : \Lambda_{P_1} \to \Lambda_{P_2}$ induced by f_P sends $\mathcal{L}_1$ isomorphically to $\mathcal{L}_2$. $(P_1, \mathcal{L}_1)$ and $(P_2, \mathcal{L}_2)$ are *isomorphic* if there exists an isomorphism between them.

The isomorphism class of the characteristic pair $(P_X, \mathcal{L}_X)$ is an invariant of a local T^n-action on X.

Lemma 4.8. *For $i = 1, 2$, let $(X_i, \mathcal{T}_i)$ be a $2n$-dimensional manifold with a local T^n-action. If there is a C^0 isomorphism $f_X \colon (X_1, \mathcal{T}_1) \to (X_2, \mathcal{T}_2)$, then f_X induces an isomorphism $f_{P_X} \colon (P_{X_1}, \mathcal{L}_{X_1}) \to (P_{X_2}, \mathcal{L}_{X_2})$ between characteristic pairs associated with X_1 and X_2.*

Proof. Let $\{(U^{X_1}_\beta, \varphi^{X_1}_\beta)\}_{\beta \in \mathcal{B}} \in \mathcal{T}_1$ and $\{(U^{X_2}_\alpha, \varphi^{X_2}_\alpha)\}_{\alpha \in \mathcal{A}} \in \mathcal{T}_2$ be maximal weakly standard atlases of X_1 and X_2, and $\{(U^{B_1}_\beta, \varphi^{B_1}_\beta)\}_{\beta \in \mathcal{B}}$ and $\{(U^{B_2}_\alpha, \varphi^{B_2}_\alpha)\}_{\alpha \in \mathcal{A}}$ atlases of B_{X_1} and B_{X_2} induced by $\{(U^{X_1}_\beta, \varphi^{X_1}_\beta)\}_{\beta \in \mathcal{B}}$ and $\{(U^{X_2}_\alpha, \varphi^{X_2}_\alpha)\}_{\alpha \in \mathcal{A}}$, respectively. Suppose that $f_X \colon (X_1, \mathcal{T}_1) \to (X_2, \mathcal{T}_2)$ is a C^0 isomorphism and $f_B \colon B_{X_1} \to B_{X_2}$ is the homeomorphism induced by f_X. By definition, on each nonempty overlap $U^{B_1}_\beta \cap (f_B)^{-1}(U^{B_2}_\alpha)$, there exists an automorphism $\rho^f_{\alpha\beta}$ of T^n such that $\varphi^{X_2}_\alpha \circ f_X \circ (\varphi^{X_1}_\beta)^{-1}$ is $\rho^f_{\alpha\beta}$-equivariant. It is easy to see that the following equality holds

$$\rho^f_{\alpha_0,\beta_0} \circ \rho^{X_1}_{\beta_0\beta_1} = \rho^{X_2}_{\alpha_0\alpha_1} \circ \rho^f_{\alpha_1\beta_1} \tag{4.4}$$

on a nonempty intersection $U^{B_1}_{\beta_0\beta_1} \cap (f_B)^{-1}(U^{B_2}_{\alpha_0\alpha_1})$, where $\rho^{X_1}_{\beta_0\beta_1}$ and $\rho^{X_2}_{\alpha_0\alpha_1}$ are automorphisms of T^n in (ii) of Definition 1.1 with respect to X_1 and X_2, respectively. We define the bundle isomorphism $(f_P)_{\alpha\beta}\colon U^{B_1}_{\beta} \cap f_B^{-1}(U^{B_2}_{\alpha}) \times \mathrm{Aut}(T^n) \to f_B(U^{B_1}_{\beta}) \cap U^{B_2}_{\alpha} \times \mathrm{Aut}(T^n)$ by

$$(f_P)_{\alpha\beta}(b,h) := (f_B(b), \rho^{f}_{\alpha\beta} \circ h).$$

Then, (4.4) implies that the equation

$$(f_P)_{\alpha_0\beta_0} \circ \varphi^{P_{X_1}}_{\beta_0\beta_1} = \varphi^{P_{X_2}}_{\alpha_0\alpha_1} \circ (f_P)_{\alpha_1\beta_1}$$

holds on $U^{B_1}_{\beta_0\beta_1} \cap f_B^{-1}(U^{B_2}_{\alpha_0\alpha_1})$, where $\varphi^{P_{X_1}}_{\beta_0\beta_1}$ (resp. $\varphi^{P_{X_2}}_{\alpha_0\alpha_1}$) is the overlap map defined by (4.2) for $\pi_{P_{X_1}}: P_{X_1} \to B_{X_1}$ (resp. $\pi_{P_{X_2}}: P_{X_2} \to B_{X_2}$) on $U^{B_1}_{\beta_0\beta_1}$ (resp. $U^{B_2}_{\alpha_0\alpha_1}$). Therefore, we can patch them together to obtain the bundle isomorphism $f_P\colon P_{X_1} \to P_{X_2}$ which covers f_B. This proves the lemma. □

4.2. **Canonical models.** In [10, Section 1.5], Davis-Januszkiewicz constructed the canonical model of a quasi-toric manifold from the based polytope and the characteristic function. A similar construction can be done by using a characteristic pair in the following way. Let B be an n-dimensional C^0 manifold with corners and $(P, \mathcal{L})$ a characteristic pair on B. We denote by $\pi_T : T_P \to B$ the T^n-bundle associated with P by the natural action of $\mathrm{Aut}(T^n)$ on T^n. First we shall explain that for any k-dimensional part $\mathcal{S}^{(k)}B$, $(P, \mathcal{L})$ determines a rank $n-k$ subtorus bundle of the restriction of $\pi_T : T_P \to B$ to $\mathcal{S}^{(k)}B$. Let $\{U_\alpha\}$ be an open covering of B such that on each U_α there exists a local trivialization $\varphi^P_\alpha\colon \pi_P^{-1}(U_\alpha) \to U_\alpha \times \mathrm{Aut}(T^n)$. On each nonempty overlap $U_{\alpha\beta}$ we denote by $\rho_{\alpha\beta}$ the transition function with respect to φ^P_α and φ^P_β, namely,

$$\varphi^P_\alpha \circ (\varphi^P_\beta)^{-1}(b, f) = (b, \rho_{\alpha\beta} f)$$

for $(b, f) \in U_\beta \times \mathrm{Aut}(T^n)$. Note that $\rho_{\alpha\beta}$ is locally constant since $\mathrm{Aut}(T^n)$ is discrete. For simplicity we assume that each $U_{\alpha\beta}$ is connected so that $\rho_{\alpha\beta}$ can be thought of as an element of $\mathrm{Aut}(T^n)$. φ^P_α induces a local trivializations of the associated bundles T_P and Λ_P which are denoted by $\varphi^T_\alpha : \pi_T^{-1}(U_\alpha) \to U_\alpha \times T^n$ and $\varphi^\Lambda_\alpha : \pi_\Lambda^{-1}(U_\alpha) \to U_\alpha \times \Lambda$, respectively. For $\mathcal{S}^{(k)}B$ we take U_α with $U_\alpha \cap \mathcal{S}^{(k)}B \neq \emptyset$. By replacing U_α by a sufficiently small one if necessary, we may assume that the intersection $U_\alpha \cap \mathcal{S}^{(n-1)}B$ has exactly $n-k$ connected components, say $(U_\alpha \cap \mathcal{S}^{(n-1)}B)_1, \cdots, (U_\alpha \cap \mathcal{S}^{(n-1)}B)_{n-k}$. For $k = n$, this means that U_α is contained in $\mathcal{S}^{(n)}B$. For $k < n$, there are $n-k$ rank one sublattices $L_1, \ldots, L_{n-k}$ of Λ such that for $a = 1, \ldots, n-k$ φ^Λ_α sends the restriction of $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ to $(U_\alpha \cap \mathcal{S}^{(n-1)}B)_a$ isomorphically to the trivial rank one subbundle $(U_\alpha \cap \mathcal{S}^{(n-1)}B)_a \times L_a$ of $(U_\alpha \cap \mathcal{S}^{(n-1)}B)_a \times \Lambda$. Since $\mathcal{L}$ is unimodular, $L_1, \ldots, L_{n-k}$ generate the $(n-k)$-dimensional subtorus of T^n which is denoted by $Z_{U_\alpha \cap \mathcal{S}^{(k)}B}$. For $k = n$, we define $Z_{U_\alpha \cap \mathcal{S}^{(n)}B}$ to be the trivial subgroup. Note that when $(P, \mathcal{L})$, $\{U_\alpha\}$, and φ^P_α are induced by some local T^n-action $\mathcal{T}$ on X, $Z_{U_\alpha \cap \mathcal{S}^{(k)}B_X}$ is the common $(n-k)$-dimensional stabilizer of T^n-action on $\mu_{\mathbb{C}^n}^{-1}(U_\alpha \cap \mathcal{S}^{(k)}B_X)$.

Suppose that another U_β satisfies the above condition and $U_{\alpha\beta} \cap \mathcal{S}^{(k)}B \neq \emptyset$. By the definition of $(P, \mathcal{L})$, $\rho_{\alpha\beta}$ sends $Z_{U_\beta \cap \mathcal{S}^{(k)}B_X}$ isomorphically to $Z_{U_\alpha \cap \mathcal{S}^{(k)}B_X}$. Hence, by patching them together by using $\rho_{\alpha\beta}$s, we obtain a rank $n-k$ subtorus bundle, which is denoted by $\pi_{Z_{\mathcal{S}^{(k)}B}}\colon Z_{\mathcal{S}^{(k)}B} \to \mathcal{S}^{(k)}B$, of the restriction of $\pi_T\colon T_P \to B$ to $\mathcal{S}^{(k)}B$.

Definition 4.9. We define two elements t, $t' \in T_P$ to be *equivalent* or $t \sim_{can} t'$ if $\pi_T(t) = \pi_T(t')$ and $t't^{-1} \in \pi^{-1}_{Z_{\mathcal{S}^{(k)}B}}(\pi_T(t))$ provided that $\pi_T(t)$ lies in $\mathcal{S}^{(k)}B$. Note that a fiber of $\pi_T \colon T_P \to B$ is equipped with the structure of a group since its structure group is $\mathrm{Aut}(T^n)$.

We denote by $X_{(P,\mathcal{L})}$ the quotient space of T_P by the equivalence relation. The bundle projection $\pi_T \colon T_P \to B$ descends to the map $\mu_{X_{(P,\mathcal{L})}} \colon X_{(P,\mathcal{L})} \to B$. Since $\mathcal{L}$ is unimodular and B is a manifold with corners, by the same way as in Davis-Januszkiewicz [10, Section 1.5], or Masuda-Panov [27, Section 3.2], we can show that $X_{(P,\mathcal{L})}$ is equipped with a C^0 local T^n-action whose orbit space is B and whose orbit map is $\mu_{X_{(P,\mathcal{L})}}$.

Definition 4.10. We call $X_{(P,\mathcal{L})}$ the *canonical model* associated with $(P, \mathcal{L})$. In particular, when $(P, \mathcal{L})$ is the characteristic pair $(P_X, \mathcal{L}_X)$ of a local T^n-action $\mathcal{T}$ on a $2n$-dimensional manifold X, we also call $X_{(P_X,\mathcal{L}_X)}$ the canonical model associated with $(X, \mathcal{T})$.

Note that by the construction, a fiber of $\mu_{X_{(P,\mathcal{L})}} \colon X_{(P,\mathcal{L})} \to B$ admits a group structure.

We give properties of a canonical model.

Proposition 4.11. *For a characteristic pair $(P, \mathcal{L})$, $\mu_{X_{(P,\mathcal{L})}} \colon X_{(P,\mathcal{L})} \to B$ admits a C^0 section.*

Proof. Since a fiber of $\pi_T \colon T_P \to B$ admits a structure of a group, it admits the section which assigns to an element $b \in B$ the unit element of $\pi_T^{-1}(b)$. The composition of this section and the natural projection from T_P to $X_{(P,\mathcal{L})}$ defines the C^0 section s of $\mu_{X_{(P,\mathcal{L})}} \colon X_{(P,\mathcal{L})} \to B$. $\square$

Proposition 4.12. *For a $2n$-dimensional manifold $(X, \mathcal{T})$ equipped with a local T^n-action, we denote the associated T^n-bundle T_{P_X} of P_X by $\pi_{T_X} \colon T_X \to B_X$ for simplicity. Then T_X acts fiberwise on X. Similarly $X_{(P_X,\mathcal{L}_X)}$ also acts fiberwise on X. For any $b \in B_X$ the action of $\mu^{-1}_{X_{(P,\mathcal{L})}}(b)$ on $\mu_X^{-1}(b)$ is simply transitive.*

Proof. Let $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ be a weakly standard atlas of X and $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ the atlas of B_X induced by $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$. For $b \in B_X$ which lies in U_α^B, the action of $\pi_{T_X}^{-1}(b)$ on $\mu_X^{-1}(b)$ can be written by

$$t \cdot x := (\varphi_\alpha^X)^{-1}(t_\alpha \cdot \varphi_\alpha^X(x))$$

for $t \in \pi_{T_X}^{-1}(b)$ and $x \in \mu_X^{-1}(b)$, where $\varphi_\alpha^T(t) = (b, t_\alpha)$. Recall that $\varphi_\alpha^T \colon \pi_{T_X}^{-1}(U_\alpha^B) \to U_\alpha^B \times T^n$ denotes the local trivialization of T_X which is induced by the local trivialization φ_α^P of P_X. It does not depend on the choice of U_α^B. In fact, suppose that b also lies in U_β^B and $\varphi_\beta^T(t) = (b, t_\beta)$. Since the transition function of φ_α^T and φ_β^T is $\rho_{\alpha\beta}$, we have $t_\alpha = \rho_{\alpha\beta}(t_\beta)$. By using the $\rho_{\alpha\beta}$-equivariantness of $\varphi_{\alpha\beta}^X$,

$$\begin{aligned}(\varphi_\alpha^X)^{-1}(t_\alpha \cdot \varphi_\alpha^X(x)) &= (\varphi_\beta^X)^{-1} \circ \varphi_{\beta\alpha}^X(\rho_{\alpha\beta}(t_\beta) \cdot \varphi_{\alpha\beta}^X \circ \varphi_\beta^X(x)) \\ &= (\varphi_\beta^X)^{-1}(t_\beta \cdot \varphi_\beta^X(x)).\end{aligned}$$

By Proposition 2.1 and the construction of $X_{(P_X,\mathcal{L}_X)}$, for each $b \in B_X$ the action of $\pi_{T_X}^{-1}(b)$ on $\mu_X^{-1}(b)$ descends to the action of $\mu^{-1}_{X_{(P_X,\mathcal{L}_X)}}(b)$ on $\mu_X^{-1}(b)$ which is simply transitive. $\square$

Lemma 4.13. *For $i = 1, 2$, let B_i be an n-dimensional C^0 manifold with corners and $(P_i, \mathcal{L}_i)$ a characteristic pair on B_i. Then, any isomorphism $f_P \colon (P_1, \mathcal{L}_1) \to (P_2, \mathcal{L}_2)$ induces the C^0 isomorphism $f_{X_{(P,\mathcal{L})}} \colon X_{(P_1,\mathcal{L}_1)} \to X_{(P_2,\mathcal{L}_2)}$ between canonical models of $(P_1, \mathcal{L}_1)$ and $(P_2, \mathcal{L}_2)$.*

Proof. It follows directly from the construction of a canonical model. $\square$

Remark 4.14. If there is an isomorphism $f_P\colon (P_1, \mathcal{L}_1) \to (P_2, \mathcal{L}_2)$ between characteristic pairs, then the induced C^0 isomorphism $f_{X_{(P,\mathcal{L})}}\colon X_{(P_1,\mathcal{L}_1)} \to X_{(P_2,\mathcal{L}_2)}$ is a fiberwise group isomorphism.

5. On sections of orbit maps

Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a local T^n-action.

Proposition 5.1. *If $\mu_X\colon X \to B_X$ has a C^0 section $s\colon B_X \to X$, then there exists a weakly standard C^0 atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ such that for each α the following diagram commutes*

$$\begin{tikzcd}
\mu_X^{-1}(U_\alpha^B) \arrow[r, "\varphi_\alpha^X"] & \mu_{\mathbb{C}^n}^{-1}(\varphi_\alpha^B(U_\alpha^B)) \\
U_\alpha^B \arrow[r, "\varphi_\alpha^B"] \arrow[u, "s"] & \varphi_\alpha^B(U_\alpha^B), \arrow[u, "\iota"']
\end{tikzcd}$$

where $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ is the atlas of B_X induced by $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ which satisfies the property in Remark 2.5 *and ι is the section defined by* (2.2).

Proof. Let $\{(U_\alpha^X, \psi_\alpha^X)\} \in \mathcal{T}$ be a weakly standard C^0 atlas of X and $\{(U_\alpha^B, \psi_\alpha^B)\} \in \mathcal{T}$ the atlas of B_X induced by $\{(U_\alpha^X, \psi_\alpha^X)\}$ which satisfies the property in Remark 2.5. For each α the equality

$$\theta_\alpha(b) \cdot s(b) = (\psi_\alpha^X)^{-1} \circ \iota \circ \psi_\alpha^B(b)$$

for $b \in U_\alpha^B$ determines a unique local section θ_α of $\mu_{X_{(P_X,\mathcal{L}_X)}}\colon X_{(P_X,\mathcal{L}_X)} \to B_X$ on U_α^B. Now we define a new local coordinate system φ_α^X on $\mu_X^{-1}(U_\alpha^B)$ by

$$\varphi_\alpha^X(x) := \psi_\alpha^X(\theta_\alpha(\mu_X(x)) \cdot x)$$

for $x \in \mu_X^{-1}(U_\alpha^B)$. Then $\{(U_\alpha^X, \varphi_\alpha^X)\}$ is the required weakly standard atlas. □

By Proposition 2.7, the restriction $\mu_X|_{\mathcal{S}^{(n)}B_X}\colon \mu_X^{-1}(\mathcal{S}^{(n)}B_X) \to \mathcal{S}^{(n)}B_X$ of μ_X to $\mathcal{S}^{(n)}B_X$ of B_X is a T^n-bundle with structure group $T^n \rtimes \mathrm{Aut}(T^n)$. This proposition implies that the structure group can be reduced to $\mathrm{Aut}(T^n)$ if μ_X admits a section.

The following proposition shows that a section of the orbit map is unique up to C^0 isomorphisms.

Proposition 5.2. *Suppose that $\mu_X\colon X \to B_X$ has two sections s_1 and s_2. Then there exists a C^0 isomorphism f_X of X such that f_X covers the identity on B_X and preserves sections, namely, $f_X \circ s_1 = s_2$.*

Proof. Since the fiberwise action of $X_{(P_X,\mathcal{L}_X)}$ on X is simply transitive, there exists a unique C^0 section θ of $\mu_{X_{(P_X,\mathcal{L}_X)}}$ such that

$$\theta(b) \cdot s_1(b) = s_2(b).$$

Then the required C^0 isomorphism f_X of X can be obtained by

$$f_X(x) := \theta(\mu_X(x)) \cdot x.$$

□

The following lemma is a straightforward generalization of the result [10, Proposition 1.8] by Davis-Januszkiewicz and the result [27, Lemma 3.6] by Masuda-Panov.

Lemma 5.3. *$\mu_X\colon X \to B_X$ is equipped with a C^0 section if and only if there exists a C^0 isomorphism between $(X, \mathcal{T})$ and $X_{(P_X,\mathcal{L}_X)}$ which covers identity on B_X.*

Proof. The if part follows from Proposition 4.11. Conversely suppose that $\mu_X : X \to B_X$ has a section which is denoted by s. Define the surjective map $\nu : T_X \to X$ by

$$\nu(t) := t \cdot s(\pi_{T_X}(t)). \tag{5.1}$$

It is easy to see from the construction of $X_{(P_X,\mathcal{L}_X)}$ that $\nu\colon T_X \to X$ descends to the required C^0 isomorphism from $X_{(P_X,\mathcal{L}_X)}$ to X. □

Next, we investigate when $\mu_X\colon X \to B_X$ has a section. We assume that the index set $\mathcal{A}$ of the weakly standard atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ is countable ordered. By the construction of $X_{(P_X,\mathcal{L}_X)}$, there exists a C^0 isomorphism $h_\alpha\colon \mu_X^{-1}(U_\alpha^B) \to \mu_{X_{(P_X,\mathcal{L}_X)}}^{-1}(U_\alpha^B)$ covering the identity on each U_α^B such that h_α is equivariant with respect to the fiberwise action of T_X and $X_{(P_X,\mathcal{L}_X)}$. By Proposition 4.12, on each nonempty overlap $U_{\alpha\beta}^B$ the equation

$$h_\alpha \circ h_\beta^{-1}(x) = \theta_{\alpha\beta}^X(b)x \tag{5.2}$$

for $b \in U_{\alpha\beta}^B$ and $x \in \mu_{X_{(P_X,\mathcal{L}_X)}}^{-1}(b)$ determines a unique local section $\theta_{\alpha\beta}^X$ of $\mu_{X_{(P_X,\mathcal{L}_X)}} : X_{(P_X,\mathcal{L}_X)} \to B_X$ on $U_{\alpha\beta}^B$. Let $\mathscr{S}_{(P_X,\mathcal{L}_X)}$ denote the sheaf of C^0 sections of $\mu_{X_{(P_X,\mathcal{L}_X)}} : X_{(P_X,\mathcal{L}_X)} \to B_X$. Then local sections $\theta_{\alpha\beta}^X$ form a Čech one-chain $\{\theta_{\alpha\beta}^X\}$ on $\{U_\alpha^B\}$ with values in $\mathscr{S}_{(P_X,\mathcal{L}_X)}$.

Lemma 5.4. $\{\theta_{\alpha\beta}^X\}$ *is a cocycle.*

Proof. It is clear by (5.2) and Proposition 4.12. □

Let $H^1(B_X;\mathscr{S}_{(P_X,\mathcal{L}_X)})$ denote the first Čech cohomology group of B_X with values in $\mathscr{S}_{(P_X,\mathcal{L}_X)}$. By the above lemma, $\{\theta_{\alpha\beta}^X\}$ defines a cohomology class in $H^1(B_X;\mathscr{S}_{(P_X,\mathcal{L}_X)})$. We denote it by $e(X,\mathcal{T})$. It is easy to see that $e(X,\mathcal{T})$ does not depend on the choice of h_αs and depends only on the local T^n-action on X.

Definition 5.5. We call $e(X,\mathcal{T})$ the *Euler class of* $\mu_X\colon X \to B_X$.

Theorem 5.6. $\mu_X : X \to B_X$ *has a section if and only if* $e(X,\mathcal{T})$ *vanishes.*

Proof. By the standard argument $(X,\mathcal{T})$ is C^0 isomorphic to $X_{(P_X,\mathcal{L}_X)}$ which covers the identity on B_X if and only if $e(X,\mathcal{T})$ vanishes. Then the theorem follows directly from this fact together with Lemma 5.3. □

Example 5.7. For an effective C^∞ T^2-action on a 4-dimensional C^∞ manifold X without nontrivial finite stabilizers, Orlik-Raymond showed in [31] that the orbit map is equipped with a section. Thus, $e(X,\mathcal{T})$ vanishes.

Example 5.8. In the case of a nonsingular toric variety, $e(X,\mathcal{T})$ vanishes. In fact, the system of standard coordinate neighborhoods induced by top-dimensional cones has the property in Proposition 5.1. Similarly, $e(X,\mathcal{T})$ vanishes for a quasi-toric manifold. It follows from [10, Lemma 1.4].

Example 5.9. If the local T^n-action is induced by a locally standard T^n-action and $\partial B_X = \emptyset$, then $\mu_X\colon X \to B_X$ is a principal T^n-bundle. In this case, $e(X,\mathcal{T})$ is nothing but the Euler class of the principal T^n-bundle.

For $i = 1, 2$, let B_i be an n-dimensional C^0 manifold with corners and $(P_i,\mathcal{L}_i)$ a characteristic pair on B_i. Suppose that there exists an isomorphism $f_P\colon (P_1,\mathcal{L}_1) \to (P_2,\mathcal{L}_2)$. By Lemma 4.13, it induces the isomorphism $f_P^*\colon H^1(B_2;\mathscr{S}_{(P_2,\mathcal{L}_2)}) \to H^1(B_1;\mathscr{S}_{(P_1,\mathcal{L}_1)})$ between cohomology groups. In particular, by Lemma 4.8 and Lemma 4.13, a C^0 isomorphism $f_X\colon (X_1,\mathcal{T}_1) \to (X_2,\mathcal{T}_2)$ induces an isomorphism $f_{P_X}^*\colon H^1(B_{X_2};\mathscr{S}_{(P_{X_2},\mathcal{L}_{X_2})}) \to H^1(B_{X_1};\mathscr{S}_{(P_{X_1},\mathcal{L}_{X_1})})$.

Lemma 5.10. *For $i = 1, 2$, let $(X_i, \mathcal{T}_i)$ be a $2n$-dimensional manifold equipped with a local T^n-action. If there is a C^0 isomorphism $f_X\colon X_1 \to X_2$, then $f^*_{P_X} e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$.*

6. The topological classification

The following is the main theorem of this paper.

Theorem 6.1. *For $i = 1, 2$, let $(X_i, \mathcal{T}_i)$ be a $2n$-dimensional manifold with a local T^n-action. $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ are C^0 isomorphic if and only if there exists an isomorphism $f_P\colon (P_{X_1}, \mathcal{L}_{X_1}) \to (P_{X_2}, \mathcal{L}_{X_2})$ between characteristic pairs associated with $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ such that $f^*_P e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$. Moreover, for any characteristic pair $(P, \mathcal{L})$ on an n-dimensional C^0 manifold B with corners and for any cohomology class $e \in H^1(B; \mathscr{S}_{(P,\mathcal{L})})$, there exists a $2n$-dimensional C^0 manifold $(X, \mathcal{T})$ equipped with a C^0 local T^n-action whose characteristic pair and the Euler class of the orbit map are equal to $(P, \mathcal{L})$ and e, respectively.*

Proof. If there exists a C^0 isomorphism $f_X : (X_1, \mathcal{T}_1) \to (X_2, \mathcal{T}_2)$, then, by Lemma 4.8 and Lemma 5.10, f_X induces an isomorphism $f_{P_X} : (P_{X_1}, \mathcal{L}_{X_1}) \to (P_{X_2}, \mathcal{L}_{X_2})$ such that $f^*_{P_X} e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$.

Conversely, suppose that there is an isomorphism $f_P\colon (P_{X_1}, \mathcal{L}_{X_1}) \to (P_{X_2}, \mathcal{L}_{X_2})$ such that $f^*_P e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$. We denote by $f_B\colon B_{X_1} \to B_{X_2}$ the stratification preserving homeomorphism which is induced by f_P. Let $\{(U^{X_1}_\beta, \varphi^{X_1}_\beta)\} \in \mathcal{T}_1$, $\{(U^{X_2}_\alpha, \varphi^{X_2}_\alpha)\} \in \mathcal{T}_2$ be weakly standard atlases of X_1, X_2 and $\{(U^{B_1}_\beta, \varphi^{B_1}_\beta)\}$, $\{(U^{B_2}_\alpha, \varphi^{B_2}_\alpha)\}$ the atlases of B_{X_1}, B_{X_2} induced by $\{(U^{X_1}_\beta, \varphi^{X_1}_\beta)\}$, $\{(U^{X_2}_\alpha, \varphi^{X_2}_\alpha)\}$ such that for each α, β, we can take C^0 isomorphisms $h^1_\beta\colon \mu^{-1}_{X_1}(U^{B_1}_\beta) \to \mu^{-1}_{X_{(P_{X_1}, \mathcal{L}_{X_1})}}(U^{B_1}_\beta)$, $h^2_\alpha\colon \mu^{-1}_{X_2}(U^{B_2}_\alpha) \to \mu^{-1}_{X_{(P_{X_2}, \mathcal{L}_{X_2})}}(U^{B_2}_\alpha)$ which satisfy (5.2) for X_1, X_2, respectively. By the assumption $f^*_P e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$, by replacing $\{(U^{X_1}_\beta, \varphi^{X_1}_\beta)\}$ by a refinement if necessary, we may assume that there exists a Čech zero-cochain $\{\theta_\beta\}$ on $\{(U^{B_1}_\beta, \varphi^{B_1}_\beta)\}$ with values in $\mathscr{S}_{X_{(P_{X_1}, \mathcal{L}_{X_1})}}$ such that

$$f^{-1}_{X_{(P,\mathcal{L})}} \circ \theta^{X_2}_{\beta_0\beta_1} \circ f_B(b) = \theta^{X_1}_{\beta_0\beta_1}(b)\theta_{\beta_1}(b)\theta_{\beta_0}(b)^{-1} \tag{6.1}$$

for $b \in U^{B_1}_{\beta_0\beta_1}$. For each nonempty overlap $U^{B_1}_\beta \cap f^{-1}_B(U^{B_2}_\alpha)$ we define the C^0 isomorphism $f_{\alpha\beta}\colon \mu^{-1}_{X_1}(U^{B_1}_\beta \cap f^{-1}_B(U^{B_2}_\alpha)) \to \mu^{-1}_{X_2}(f_B(U^{B_1}_\beta) \cap U^{B_2}_\alpha)$ by

$$f_{\alpha\beta}(x) := (h^2_\alpha)^{-1} \circ f_{X_{(P,\mathcal{L})}} \left(\theta_\beta(\mu_{X_1}(x)) h^1_\beta(x)\right).$$

By using (6.1) and (5.2) for X_1 and X_2, for any $x \in \mu^{-1}_{X_1}(U^{B_1}_{\beta_0\beta_1} \cap f^{-1}_B(U^{B_2}_{\alpha_0\alpha_1}))$ with $b = \mu_{X_1}(x)$

$$\begin{aligned}
f_{\alpha_0\beta_0}(x) =& (h^2_{\alpha_0})^{-1} \circ f_{X_{(P,\mathcal{L})}} \left(\theta_{\beta_0}(b) h^1_{\beta_0}(x)\right) \\
&= (h^2_{\alpha_1})^{-1} \left(\theta^{X_2}_{\alpha_0\alpha_1}(f_B(b)) f_{X_{(P,\mathcal{L})}} \left(\theta_{\beta_0}(b)\theta^{X_1}_{\beta_0\beta_1}(b)^{-1} h^1_{\beta_1}(x)\right)\right) \\
&= (h^2_{\alpha_1})^{-1} \left(\theta^{X_2}_{\alpha_0\alpha_1}(f_B(b)) f_{X_{(P,\mathcal{L})}} \left(\theta_{\beta_0}(b)\theta^{X_1}_{\beta_0\beta_1}(b)^{-1}\right) f_{X_{(P,\mathcal{L})}} \left(h^1_{\beta_1}(x)\right)\right) \\
&= (h^2_{\alpha_1})^{-1} \left(f_{X_{(P,\mathcal{L})}} \left(\theta_{\beta_1}(b)\right) f_{X_{(P,\mathcal{L})}} \left(h^1_{\beta_1}(x)\right)\right) \\
&= (h^2_{\alpha_1})^{-1} \circ f_{X_{(P,\mathcal{L})}} \left(\theta_{\beta_1}(b) h^1_{\beta_1}(x)\right) \\
&= f_{\alpha_1\beta_1}(x),
\end{aligned}$$

where we used the fact that $f_{X_{(P,\mathcal{L})}} : X_{(P_1,\mathcal{L}_1)} \to X_{(P_2,\mathcal{L}_2)}$ is fiberwise group isomorphism (see Remark 4.14). Then we can patch the local C^0 isomorphisms $f_{\alpha\beta}$ together to obtain the required C^0 isomorphism f_X.

Suppose that $(P, \mathcal{L})$ is a characteristic pair on an n-dimensional C^0 manifold B with corners and e is an element of $H^1(B; \mathscr{S}_{(P,\mathcal{L})})$. We take a representative $\{\theta_{\alpha\beta}\}$ of e on a sufficiently small open cover $\{U_\alpha\}$ of B. Then we can construct a new $2n$-dimensional C^0 manifold $(X, \mathcal{T})$ equipped with a C^0 local T^n-action by setting

$$X := \left(\coprod_{\alpha} \mu^{-1}_{X_{(P,\mathcal{L})}}(U^B_\alpha) \right) / \sim,$$

where $x_\alpha \in \mu^{-1}_{X_{(P,\mathcal{L})}}(U^B_\alpha) \sim x_\beta \in \mu^{-1}_{X_{(P,\mathcal{L})}}(U^B_\beta)$ if and only if $\mu_{X_{(P,\mathcal{L})}}(x_\alpha) = \mu_{X_{(P,\mathcal{L})}}(x_\beta)$ and $x_\alpha = \theta_{\alpha\beta}(\mu_{X_{(P,\mathcal{L})}}(x_\alpha))x_\beta$. It is easy to see that its characteristic pair and Euler class of the orbit map are equal to $(P, \mathcal{L})$ and e, respectively. □

Next we focus on the case of locally standard torus actions. We take notice that if a manifold X is equipped with a locally standard torus action, then, P_X is the product bundle $P_X = B_X \times \mathrm{Aut}(T^n)$. In this case, we can obtain the following theorem.

Theorem 6.2. *For $i = 1, 2$, let X_i be a $2n$-dimensional manifold equipped with a locally standard T^n-action and ρ an automorphism of T^n. X_1 and X_2 are ρ-equivariant homeomorphic if and only if there exists a stratification preserving homeomorphism $f_B \colon B_{X_1} \to B_{X_2}$ such that the bundle isomorphism $f_P \colon P_{X_1} \to P_{X_2}$ defined by $f_P := f_B \times \rho$ sends $\mathcal{L}_{X_1}$ isomorphically to $\mathcal{L}_{X_2}$ and $f_P^* e(X_2, \mathcal{T}_2) = e(X_1, \mathcal{T}_1)$. Moreover, for any characteristic pair $(P, \mathcal{L})$, where P is trivial, on an n-dimensional C^0 manifold B with corners and for any cohomology class $e \in H^1(B; \mathscr{S}_{(P,\mathcal{L})})$, there exists a $2n$-dimensional C^0 manifold equipped with a locally standard T^n-action whose characteristic pair and the Euler class of the orbit map are equal to $(P, \mathcal{L})$ and e, respectively.*

The proof is almost same as that of Theorem 6.1. In particular, by putting $\rho = id_{T^n}$, we obtain the following corollary.

Corollary 6.3. *Locally standard torus actions are classified by the characteristic bundle and the Euler class of the orbit map up to equivariant homeomorphisms.*

7. The symplectic case

In this section we identify Λ with $\mathbb{Z}^n$ by using the fixed decomposition $T^n = (S^1)^n$. By using the natural isomorphism $\mathrm{Aut}(T^n) \to \mathrm{GL}(\Lambda)$ together with this fact, we also identify $\mathrm{Aut}(T^n)$ with $\mathrm{GL}_n(\mathbb{Z})$.

7.1. **Integral affine structures.**

Let B be an n-dimensional C^∞ manifold with corners and $\{(U^B_\alpha, \varphi^B_\alpha)\}$ an atlas of B consisting of coordinate neighborhoods modeled on open subsets of $\mathbb{R}^n_+$.

Definition 7.1. $\{(U^B_\alpha, \varphi^B_\alpha)\}$ is called an *integral affine structure* on B if on each nonempty overlap $U^B_{\alpha\beta} \neq \emptyset$, there exists an element $A_{\alpha\beta}$ of $\mathrm{GL}_n(\mathbb{Z})$ and there also exists a constant $c_{\alpha\beta} \in \mathbb{R}^n$ such that the overlap map $\varphi^B_{\alpha\beta} \colon \varphi^B_\beta(U^B_{\alpha\beta}) \to \varphi^B_\alpha(U^B_{\alpha\beta})$ is of the form

$$\varphi^B_{\alpha\beta}(\xi) = A_{\alpha\beta}(\xi) + c_{\alpha\beta}. \tag{7.1}$$

First we show the following lemma.

Lemma 7.2. *For an n-dimensional C^∞ manifold B equipped with an integral affine structure $\{(U^B_\alpha, \varphi^B_\alpha)\}$, there is a characteristic pair on B associated with $\{(U^B_\alpha, \varphi^B_\alpha)\}$.*

Proof. By definition, the structure group of the cotangent bundle T^*B of B is reduced to $\mathrm{GL}_n(\mathbb{Z})$. We denote by $\pi_P\colon P \to B$ the frame bundle of T^*B with structure group $\mathrm{GL}_n(\mathbb{Z})$ and also denote by $\pi_\Lambda\colon \Lambda_P \to B$ the associated Λ-bundle of $\pi_P\colon P \to B$. Let $(U^B_\alpha, \varphi^B_\alpha)$ be a coordinate neighborhood in $\{(U^B_\alpha, \varphi^B_\alpha)\}$ with $U^B_\alpha \cap \mathcal{S}^{(n-1)}B_X \neq \emptyset$. We may assume that the intersection $U^B_\alpha \cap \mathcal{S}^{(n-1)}B_X$ is connected. (Otherwise, we may consider componentwise.) Then there exists a unique primitive vector $u_\alpha \in \Lambda$ such that $\varphi^B_\alpha(U^B_\alpha)$ and $\varphi^B_\alpha(U^B_\alpha \cap \mathcal{S}^{(n-1)}B_X)$ are contained in the upper half space $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\alpha\rangle \geq 0\}$ and the hyperplane $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\alpha\rangle = 0\}$ determined by u_α, respectively. Suppose that $(U^B_\beta, \varphi^B_\beta)$ is another coordinate neighborhoods satisfying the above conditions and the intersection $U^B_{\alpha\beta}\cap \mathcal{S}^{(n-1)}B_X$ is nonempty. Let $u_\beta \in \Lambda$ be the corresponding primitive vector. Since the overlap map $\varphi^B_{\alpha\beta}$ is of the form (7.1) and since $\varphi^B_{\alpha\beta}$ sends $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\beta\rangle \geq 0\}$ and $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\beta\rangle = 0\}$ diffeomorphically to $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\alpha\rangle \geq 0\}$ and $\{\xi \in \mathbb{R}^n\colon \langle \xi, u_\alpha\rangle = 0\}$, respectively, we can show that $u_\alpha = A^{-T}_{\alpha\beta}(u_\beta)$. In particular, u_α's form a section s of $\pi_\Lambda|_{\mathcal{S}^{(n-1)}B}\colon \Lambda_P|_{\mathcal{S}^{(n-1)}B} \to \mathcal{S}^{(n-1)}B$. Let $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ be the rank one sublattice bundle of $\pi_\Lambda|_{\mathcal{S}^{(n-1)}B}\colon \Lambda_P|_{\mathcal{S}^{(n-1)}B} \to \mathcal{S}^{(n-1)}B$ which is fiberwise generated by s. By construction, $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ is unimodular, hence the pair $(P, \mathcal{L})$ is a characteristic pair on B. ☐

Note that, by the proof of this lemma, the characteristic bundle $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ admits a section which generates $\pi_{\mathcal{L}}\colon \mathcal{L} \to \mathcal{S}^{(n-1)}B$ fiberwise.

In Section 4.2 we constructed the canonical model from a characteristic pair. Next we show that the canonical model constructed from the characteristic pair associated with an integral affine structure is a C^∞ manifold and admits a symplectic structure such that the orbit map is a locally toric Lagrangian fibration. Let B be an n-dimensional C^∞ manifold equipped with an integral affine structure $\{(U^B_\alpha, \varphi^B_\alpha)\}$ and $(P, \mathcal{L})$ the characteristic pair associated with $\{(U^B_\alpha, \varphi^B_\alpha)\}$. Let $\pi_T\colon T_P \to B$ be the associated T^n-bundle of $\pi_P\colon P \to B$. Then we have the following exact sequence of associated fiber bundles of P

$$\begin{tikzcd} 0 \arrow[r] & \Lambda_P \arrow[r] & T^*B \arrow[r] & T_P \arrow[r] & 0. \end{tikzcd}$$

As is well-known, T^*B is equipped with the standard symplectic structure ω_{T^*B}. Since the natural fiberwise action of Λ_P on T^*B preserves ω_{T^*B}, ω_{T^*B} descends to a symplectic structure on T_P, which is denoted by ω_{T_P}, so that $\pi_T\colon (T_P, \omega_{T_P}) \to B$ is a nonsingular Lagrangian fibration. Moreover, we can show that following lemma.

Lemma 7.3. *The canonical model $X_{(P,\mathcal{L})}$ is a C^∞ manifold and admits a symplectic structure $\omega_{X_{(P,\mathcal{L})}}$ so that $\mu_{X_{(P,\mathcal{L})}}\colon (X_{(P,\mathcal{L})}, \omega_{X_{(P,\mathcal{L})}}) \to B$ a locally toric Lagrangian fibration.*

Proof. In this case $X_{(P,\mathcal{L})}$ can be reconstructed from (T_P, ω_{T_P}) by using the symplectic cutting technique in the following way. For each $(U^B_\alpha, \varphi^B_\alpha)$, $\overline{\varphi}^T_\alpha\colon \pi_T^{-1}(U^B_\alpha) \to \varphi^B_\alpha(U^B_\alpha) \times T^n$ denotes the composition of $\varphi^B_\alpha \times \mathrm{id}_{T^n}\colon U^B_\alpha \times T^n \to \varphi^B_\alpha(U^B_\alpha) \times T^n$ and the local trivialization φ^T_α of T_P on U^B_α induced by the local trivialization φ^P_α of P on U^B_α. By the construction of ω_{T_P}, $\overline{\varphi}^T_\alpha$ preserves the symplectic structures, namely, $(\overline{\varphi}^T_\alpha)^*\omega_{\mathbb{R}^n\times T^n} = \omega_{T_P}$. We put

$$I_\alpha := \{i \in \{1, \dots, n\}\colon \varphi^B_\alpha(U^B_\alpha) \cap \{\xi \in \mathbb{R}^n_+\colon \xi_i = 0\} \neq \emptyset\}$$

and

$$T_{I_\alpha} := \{u \in T^n\colon u_i = 1 \text{ unless } i \in I_\alpha\}.$$

Suppose that $I_\alpha = \{i_1, \dots, i_k\}$ and $i_1 < \dots < i_k$. We take an open set V_α of $\mathbb{R}^n$ which satisfies $\varphi^B_\alpha(U^B_\alpha) = V_\alpha \cap \mathbb{R}^n_+$. T_{I_α} acts on $V_\alpha \times T^n \times \mathbb{C}^k$ by

$$u \cdot (\xi, v, z) := \left(\xi, uv, (u^{-1}_{i_1} z_1, \dots, u^{-1}_{i_k} z_k)\right).$$

This action is Hamiltonian with respect to the symplectic structure $\omega_{\mathbb{R}^n\times T^n}\oplus\omega_{\mathbb{C}^k}$ and a moment map $\Phi_\alpha\colon (V_\alpha\times T^n\times\mathbb{C}^k, \omega_{\mathbb{R}^n\times T^n}\oplus\omega_{\mathbb{C}^k})\to\mathbb{R}^k$ is written by

$$\Phi_\alpha(\xi,v,z)=(\xi_{i_1}-|z_1|^2,\ldots,\xi_{i_k}-|z_k|^2).$$

We take the symplectic quotient $\Phi_\alpha^{-1}(0)/T_{I_\alpha}$ of the T_{I_α}-action at $0\in\mathbb{R}^k$. We denote by ω_α the reduced symplectic structure on $\Phi_\alpha^{-1}(0)/T_{I_\alpha}$. The natural action of T^n on $V_\alpha\times T^n\times\mathbb{C}^k$ by the multiplication of the second factor induces a Hamiltonian T^n-action on $(\Phi_\alpha^{-1}(0)/T_{I_\alpha},\omega_\alpha)$ with moment map $\mu_\alpha([\xi,u,z])=\xi$. Note that the image of μ_α is equal to $\varphi_\alpha^B(U_\alpha^B)$. It is obvious that this T^n-action on $(\Phi_\alpha^{-1}(0)/T_{I_\alpha},\omega_\alpha)$ with moment map μ_α is naturally identified with the standard T^n-action on $(\mu_{\mathbb{C}^n}^{-1}(\varphi_\alpha^B(U_\alpha^B)),\omega_{\mathbb{C}^n})$ with moment map $\mu_{\mathbb{C}^n}$. By the construction of $X_{(P,\mathcal{L})}$ we can also show that $\overline{\varphi}_\alpha^T$ induces a natural homeomorphism $\varphi_\alpha^{X_{(P,\mathcal{L})}}\colon \mu_{X_{(P,\mathcal{L})}}^{-1}(U_\alpha^B)\to\Phi_\alpha^{-1}(0)/T_{I_\alpha}$ which satisfies $\mu_\alpha\circ\varphi_\alpha^{X_{(P,\mathcal{L})}}=\varphi_\alpha^B\circ\mu_{X_{(P,\mathcal{L})}}$.

Suppose that $(U_\alpha^B,\varphi_\alpha^B)$ and $(U_\beta^B,\varphi_\beta^B)$ are coordinate neighborhoods with nonempty overlap $U_{\alpha\beta}^B$. To prove the lemma, it is sufficient to show that the overlap map $\varphi_{\alpha\beta}^{X_{(P,\mathcal{L})}}:=\varphi_\alpha^{X_{(P,\mathcal{L})}}\circ(\varphi_\beta^{X_{(P,\mathcal{L})}})^{-1}$ is a symplectomorphism from $(\mu_\beta^{-1}(\varphi_\beta^B(U_{\alpha\beta}^B)),\omega_\beta)$ to $(\mu_\alpha^{-1}(\varphi_\alpha^B(U_{\alpha\beta}^B)),\omega_\alpha)$. Since the overlap map $\overline{\varphi}_{\alpha\beta}^T:=\overline{\varphi}_\alpha^T\circ(\overline{\varphi}_\beta^T)^{-1}$ is of the form $\overline{\varphi}_{\alpha\beta}^T(\xi,u)=(A_{\alpha\beta}(\xi)+c_{\alpha\beta},A_{\alpha\beta}^{-T}(u))$, it can be naturally extended to the symplectomorphism from $(V_\beta\cap(\varphi_{\alpha\beta}^B)^{-1}(V_\alpha)\times T^n,\omega_{\mathbb{R}^n\times T^n})$ to $(\varphi_{\alpha\beta}^B(V_\beta)\cap V_\alpha\times T^n,\omega_{\mathbb{R}^n\times T^n})$, which is also denoted by the same notation $\overline{\varphi}_{\alpha\beta}^T$. Since $\overline{\varphi}_{\alpha\beta}^T$ is $A_{\alpha\beta}^{-T}$-equivariant with respect to the actions of T_{I_β} and T_{I_α}, it descends to a symplectomorphism from $(\mu_\beta^{-1}(\varphi_\beta^B(U_{\alpha\beta}^B)),\omega_\beta)$ to $(\mu_\alpha^{-1}(\varphi_\alpha^B(U_{\alpha\beta}^B)),\omega_\alpha)$ which is nothing but $\varphi_{\alpha\beta}^{X_{(P,\mathcal{L})}}$. This proves the lemma. □

Remark 7.4. Let B_1 and B_2 be n-dimensional C^∞ manifolds equipped with integral affine structures. Suppose that there exists a diffeomorphism $f_B\colon B_1\to B_2$ which preserves the integral affine structures. Then f_B induces an isomorphism $f_P\colon (P_1,\mathcal{L}_1)\to(P_2,\mathcal{L}_2)$ between the characteristic pairs associated with the integral affine structures. Moreover, it is easy to see that the C^0 isomorphism $f_{X_{(P,\mathcal{L})}}\colon X_{(P_1,\mathcal{L}_1)}\to X_{(P_2,\mathcal{L}_2)}$ induced by f_P is a fiber-preserving symplectomorphism.

7.2. **The necessary and sufficient condition.** Let $(X,\mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^∞ local T^n-action $\mathcal{T}$. In this subsection, we investigate the condition in order that X admits a symplectic structure ω so that $\mu_X\colon(X,\omega)\to B_X$ is a locally toric Lagrangian fibration.

Definition 7.5. Let $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}\in\mathcal{T}$ be a weakly standard C^∞ atlas of X and $\{(U_\alpha^B,\varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ the atlas of B_X induced by $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$. For each nonempty overlap $U_{\alpha\beta}^X\neq\emptyset$, let $\rho_{\alpha\beta}\in\mathrm{Aut}(T^n)$ be the automorphism in (ii) of Definition 1.1 with respect to $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$. We call $\{(U_\alpha^B,\varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ an *integral affine structure compatible with* $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ if $\{(U_\alpha^B,\varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ is an integral affine structure on B_X and on each nonempty overlap $U_{\alpha\beta}^B\neq\emptyset$, the element $A_{\alpha\beta}\in\mathrm{GL}_n(\mathbb{Z})$ in (7.1) is equal to $\rho_{\alpha\beta}^{-T}$.

Lemma 7.6. *Suppose that there exists a symplectic structure ω on X and there also exists a weakly standard atlas $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}\in\mathcal{T}$ of X such that on each U_α^X, φ_α^X preserves symplectic structures, namely, $\omega={\varphi_\alpha^X}^*\omega_{\mathbb{C}^n}$. Then the atlas $\{(U_\alpha^B,\varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X induced by $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ is an integral affine structure compatible with $\{(U_\alpha^X,\varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$. In particular, B_X becomes a C^∞ manifold with corners.*

Proof. By assumption, the restriction of $\mu_X\colon (X,\omega) \to B_X$ to $B_X \setminus \partial B_X$ is a nonsingular Lagrangian fibration, and as in the proof of Proposition 3.13, we can construct a symplectomorphism $\phi_\alpha\colon (\mu_X^{-1}(U_\alpha^B \setminus \partial B_X),\omega) \to (\varphi_\alpha^B(U_\alpha^B \setminus \partial B_X) \times T^n, \omega_{\mathbb{R}^n\times T^n})$ such that $\mathrm{pr}_1 \circ \phi_\alpha = \varphi_\alpha^B \circ \mu_X$ for each α. By applying Lemma 3.12 to the overlap map $\phi_{\alpha\beta} := \phi_\alpha \circ \phi_\beta^{-1}$ on each nonempty overlap $U_{\alpha\beta}^B \setminus \partial B_X$, we can show that the overlap map $\varphi_{\alpha\beta}^B$ of the base is of the form (7.1). Since $U_{\alpha\beta}^B \setminus \partial B_X$ is open dense in $U_{\alpha\beta}^B$, $\varphi_{\alpha\beta}^B$ should be of the form (7.1) on the whole $U_{\alpha\beta}^B$. □

Let $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ be a weakly standard atlas of X. Suppose that the induced atlas $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X is an integral affine structure compatible with $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$. Then the principal $\mathrm{Aut}(T^n)$-bundle P_X is nothing but the frame bundle of T^*B_X and the characteristic pair $(P_X, \mathcal{L}_X)$ is nothing but the characteristic pair associated with $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$. By Lemma 7.3, the canonical model $X_{(P_X,\mathcal{L}_X)}$ is a C^∞ symplectic manifold with symplectic structure $\omega_{X_{(P_X,\mathcal{L}_X)}}$ so that $\mu_{X_{(P_X,\mathcal{L}_X)}}\colon (X_{(P_X,\mathcal{L}_X)}, \omega_{X_{(P_X,\mathcal{L}_X)}}) \to B_X$ a locally toric Lagrangian fibration. In particular, the local isomorphism $h_\alpha\colon \mu_X^{-1}(U_\alpha^B) \to \mu_{X_{(P_X,\mathcal{L}_X)}}^{-1}(U_\alpha^B)$ in Section 5 can be taken to be a C^∞ isomorphism which covers the identity on each U_α^B and by [18, Lemma 3.2] $\theta_{\alpha\beta}^X$ defined by (5.2) can be also taken to be a C^∞ local section of T_X on $U_{\alpha\beta}^B$. Then a necessary and sufficient condition is given as follows.

Theorem 7.7. *Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold equipped with a C^∞ local T^n-action $\mathcal{T}$. There exists a symplectic structure ω on X and there also exists a weakly standard atlas $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}} \in \mathcal{T}$ of X such that on each U_α^X, $\omega = {\varphi_\alpha^X}^*\omega_{\mathbb{C}^n}$ if and only if the atlas $\{(U_\alpha^B, \varphi_\alpha^B)\}_{\alpha\in\mathcal{A}}$ of B_X induced by $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ is an integral affine structure compatible with $\{(U_\alpha^X, \varphi_\alpha^X)\}_{\alpha\in\mathcal{A}}$ and on each nonempty overlap $U_{\alpha\beta}^B$, $\theta_{\alpha\beta}^X$ is a Lagrangian section, namely, $(\theta_{\alpha\beta}^X)^*\omega_{T_X}$ vanishes.*

For nonsingular Lagrangian fibrations, this result is obtained by Duistermaat [11]. See also [32], [29]. Recently, in [15] Gay-Symington showed the similar result for near-symplectic four-manifolds.

7.3. **The symplectic classification.** Suppose that $(X, \mathcal{T})$ is a $2n$-dimensional C^∞ manifold equipped with a C^∞ local T^n-action $\mathcal{T}$ satisfies the condition in Theorem 7.7. Then X is equipped with a symplectic structure ω so that $\mu_X\colon (X,\omega) \to B_X$ is a locally toric Lagrangian fibration, and the local sections $\theta_{\alpha\beta}^X$ define a Čech cohomology class $\lambda(X) \in H^1(B_X; \mathscr{S}_{T_X}^{Lag})$ of B_X with values in the sheaf $\mathscr{S}_{T_X}^{Lag}$ of Lagrangian sections of $\pi_{T_X}\colon (T_X, \omega_{T_X}) \to B_X$. $\lambda(X)$ is called a *Lagrangian class* for $\mu_X\colon (X,\omega) \to B_X$. Note that for two $2n$-dimensional C^∞ manifolds $(X_1, \mathcal{T}_1)$ and $(X_2, \mathcal{T}_2)$ with C^∞ local T^n-actions which satisfy the condition in Theorem 7.7, if there exists a diffeomorphism $f_B\colon B_{X_1} \to B_{X_2}$ which preserves the integral affine structures, then f_B induces an isomorphism $f_P^*\colon H^1(B_{X_2}; \mathscr{S}_{T_{X_2}}^{Lag}) \to H^1(B_{X_1}; \mathscr{S}_{T_{X_1}}^{Lag})$.

Theorem 7.8 ([5])**.** *For $i = 1,2$, let $(X_i, \mathcal{T}_i)$ be a $2n$-dimensional C^∞ manifold with a C^∞ local T^n-action which is equipped with a a symplectic structure ω_i so that $\mu_{X_i}\colon (X_i, \omega_i) \to B_{X_i}$ is a locally toric Lagrangian fibration. $\mu_{X_1}\colon (X_1,\omega_1) \to B_{X_1}$ and $\mu_{X_2}\colon (X_2,\omega_2) \to B_{X_2}$ are fiber-preserving symplectomorphic if and only if there exists a diffeomorphism $f_B\colon B_{X_1} \to B_{X_2}$ which preserves the integral affine structures such that $f_P^*\lambda(X_2) = \lambda(X_1)$. Moreover, for any n-dimensional C^∞ manifold B with corners equipped with an integral affine structure and for any cohomology class $\lambda \in H^1(B; \mathscr{S}_{T_P}^{Lag})$, there exists a locally toric Lagrangian fibration whose integral affine structure and the Lagrangian class are the given ones. Here $\mathscr{S}_{T_P}^{Lag}$ is the sheaf of Lagrangian sections of $\pi_T\colon (T_P, \omega_{T_P}) \to B$.*

The proof is similar to Theorem 6.1.

8. TOPOLOGY

Let $(X, \mathcal{T})$ be a $2n$-dimensional manifold with a local T^n-action. We assume that X is connected (hence B is also connected) and $e(X, \mathcal{T})$ vanishes. In this section, we investigate some topological invariants for X.

8.1. Fundamental groups.

In this subsection, we investigate the fundamental group of X. We fix a point b_0 in the interior $\mathcal{S}^{(n)}B_X$ of B_X and also fix points $x_0 \in {\mu_X}^{-1}(b_0)$ and $t_0 \in \pi_{T_X}^{-1}(b_0)$ which satisfy $\nu(t_0) = x_0$ as base points of X and T_X, respectively, where $\nu\colon T_X \to X$ is the map defined by (5.1). Comparing the fundamental group of T_X with that of X by using the homomorphism induced from $\nu\colon T_X \to X$, we have the following result.

Theorem 8.1. *Suppose that $\mathcal{S}^{(0)}B_X$ is nonempty. Then μ_X induces an isomorphism ${\mu_X}_*\colon \pi_1(X, x_0) \cong \pi_1(B_X, b_0)$ of fundamental groups.*

Proof. Since the structure group of $\pi_{T_X}: T_X \to B_X$ is $\mathrm{Aut}(T^n)$ it admits a section s_T. s_T defines a section s_X of μ_X by $s_X := \nu \circ s_T$. Now we have the following commutative diagram of split short exact sequences for fundamental groups

$$\begin{tikzcd}
1 \arrow[r] & \pi_1(\pi_{T_X}^{-1}(b_0), t_0) \arrow[r, "\iota_*"] \arrow[d, "\kappa"] & \pi_1(T_X, t_0) \arrow[r, "{\pi_{T_X}}_*"] \arrow[d, "\nu_*"] & \pi_1(B_X, b_0) \arrow[r] \arrow[d, "\mathrm{id}_{\pi_1(B_X, b_0)}"] & 1 & \text{(exact)} \\
1 \arrow[r] & \ker {\mu_X}_* \arrow[r] & \pi_1(X, x_0) \arrow[r, "{\mu_X}_*"] & \pi_1(B_X, b_0) \arrow[r] & 1 & \text{(exact),}
\end{tikzcd}$$

where $\iota\colon \pi_{T_X}^{-1}(b_0) \hookrightarrow T_X$ is the natural inclusion and $\kappa\colon \pi_1(\pi_{T_X}^{-1}(b_0), t_0) \to \ker {\mu_X}_*$ is the homomorphism induced by the restriction $\nu|_{\pi_{T_X}^{-1}(b_0)} : \pi_{T_X}^{-1}(b_0) \to \mu^{-1}(b_0)$ of ν to $\pi_{T_X}^{-1}(b_0)$. First we claim that κ is surjective. Note that it is equivalent to the surjectivity of ν_* since ${s_X}_* = \nu_* \circ {s_T}_*$. Since b_0 is in the interior of B_X, $\nu|_{\pi_{T_X}^{-1}(b_0)}$ is a homeomorphism which sends t_0 to x_0. Then it is sufficient to show that every element of $\ker {\mu_X}_*$ is represented by a loop in ${\mu_X}^{-1}(b_0)$. Let $\alpha \in \ker {\mu_X}_*$ and take its representative $a'\colon S^1 \to X$ with $a'(1) = x_0$. Then the map $\mu_X \circ a'\colon S^1 \to B_X$ is homotopic to the constant map b_0. If necessary, by replacing a representative of α, we can take a contractible open set U located in the interior of B_X such that the image of a' is included in ${\mu_X}^{-1}(U)$. Since U is in the interior of B_X, by Proposition 2.7, ${\mu_X}^{-1}(b_0)$ is a deformation retract of ${\mu_X}^{-1}(U)$. We take a deformation retraction $h : I \times {\mu_X}^{-1}(U) \to {\mu_X}^{-1}(U)$ which satisfies $h(0, \cdot) = id_{{\mu_X}^{-1}(U)}$ and $h(1, \cdot) : {\mu_X}^{-1}(U) \to {\mu_X}^{-1}(b_0)$, where I implies the unit interval $[0, 1]$. Then the map $f(s, u) := h(s, a'(u))$ is the homotopy which connects a' to the loop $a(u) := h(1, a'(u))$ in ${\mu_X}^{-1}(b_0)$ with $a(1) = x_0$. This proves the claim.

To prove the theorem, it is sufficient to show that κ is the constant map whose value is the unit element. By assumption, there exists a point b_1 in $\mathcal{S}^{(0)}B_X$. The fiber ${\mu_X}^{-1}(b_1)$ consists of only one point which is denoted by x_1. We take a path $\gamma\colon I \to B_X$ with $\gamma(0) = b_0$ and $\gamma(1) = b_1$. (Since B_X is a connected C^0 manifold, we can always take such a path.) Since $\pi_{T_X} : T_X \to B_X$ is a fiber bundle, there exists a continuous map $\widetilde{\gamma}\colon I \times \pi_{T_X}^{-1}(b_0) \to T_X$ which satisfies the followings

$$\pi_{T_X} \circ \widetilde{\gamma}(s, t) = \gamma(s), \quad \widetilde{\gamma}(0, t) = t$$

for $s \in I$ and $t \in \pi_{T_X}^{-1}(b_0)$. $\widetilde{\gamma}$ is unique up to homotopy. For any $\alpha \in \pi_1(\pi_{T_X}^{-1}(b_0), t_0)$ and its representative $a\colon S^1 \to \pi_{T_X}^{-1}(b_0)$ with $a(1) = t_0$, we define the map $\overline{a}\colon I \times S^1 \to X$ by $\overline{a}(s, u) := \nu(\widetilde{\gamma}(s, a(u)))$. $\overline{a}$ satisfies

$$\overline{a}(0, u) = \nu \circ a(u), \quad \overline{a}(1, u) \equiv x_1$$

for any $u \in S^1$ and $s \in I$. Then, $\overline{a}$ descends to the map from a two-dimensional closed disc D which bounds $\nu \circ a$. This implies that $\kappa(\alpha)$ is the unit element. $\square$

8.2. **Cohomology groups.** In this subsection, we give the method for computing cohomology groups of X.

Suppose that B_X is equipped with a structure of a CW complex so that each p-cell $e^{(p)}$ is contained in some $\mathcal{S}^{(k)}B_X$ of B_X. Let $B_X^{(p)}$ be the p-skeleton and $X^{(p)} := {\mu_X}^{-1}(B_X^{(p)})$ its preimage by μ_X. We put

$$A^{p,q} := H^{p+q}(X, X^{(p-1)}; \mathbb{Z}), \ E^{p,q} := H^{p+q}(X^{(p)}, X^{(p-1)}; \mathbb{Z}).$$

$i\colon A^{p,q} \to A^{p-1,q+1}$ and $j\colon A^{p,q} \to E^{p,q}$ denote the maps induced from inclusions $(X, X^{(p-2)}) \subset (X, X^{(p-1)})$ and $(X^{(p)}, X^{(p-1)}) \subset (X, X^{(p-1)})$, respectively, and $k :E^{p,q} \to A^{p+1,q}$ also denotes the connecting homomorphism of the exact sequence of the triple $(X, X^{(p)}, X^{(p-1)})$. We consider the cohomology Leray spectral sequence of $\mu_X : X \to B_X$, namely, the spectral sequence $\{(E_X)_r^{p,q}, d_r^X\}$ associated with the exact couple

$$\begin{tikzcd}
\bigoplus_{p,q} A^{p,q} \arrow[rr, "i"] & & \bigoplus_{p,q} A^{p,q} \arrow[dl, "j"] \\
& \bigoplus_{p,q} E^{p,q}. \arrow[ul, "k"] &
\end{tikzcd}$$

For the spectral sequence of cohomology groups, see [20]. For a p-cell $e^{(p)}$, we denote by $c^{(p)}$ the center of $e^{(p)}$ and also denote the restriction of $\nu : T_X \to X$ to $\pi_{T_X}^{-1}(c^{(p)})$ by $\nu_{c^{(p)}} : \pi_{T_X}^{-1}(c^{(p)}) \to {\mu_X}^{-1}(c^{(p)})$. Let $(C^*(B_X; \mathscr{H}_T^q), \delta^*)$ be the cochain complex of the CW complex B_X with the Serre local system $\mathscr{H}_T^q$ of the qth cohomology with $\mathbb{Z}$-coefficient for the fiber bundle $\pi_{T_X} : T_X \to B_X$. We denote by $C^p(B_X; \mathscr{H}_X^q)$ the subset of $C^p(B_X; \mathscr{H}_T^q)$ consisting of those cochains which take values in the image $\nu^*_{c_\lambda^{(p)}} \left(H^q({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \right)$ of $H^q({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z})$ by $\nu^*_{c_\lambda^{(p)}}$ for each p-cell $e_\lambda^{(p)}$.

Theorem 8.2. *For any q, $C^*(B_X; \mathscr{H}_X^q)$ is a subcomplex of $(C^*(B_X; \mathscr{H}_T^q), \delta)$. We denote its pth cohomology group by $H^p(B_X; \mathscr{H}_X^q)$. Then we have the isomorphisms*

$$(E_X)_1^{p,q} \cong C^p(B_X; \mathscr{H}_X^q), \quad (E_X)_2^{p,q} \cong H^p(B_X; \mathscr{H}_X^q),$$
$$(E_X)_\infty^{p,q} = F^p H^{p+q}(X; \mathbb{Z}) / F^{p+1} H^{p+q}(X; \mathbb{Z}),$$

where $F^l H^k(X; \mathbb{Z})$ is the image of the natural map $H^k(X, X^{(l-1)}; \mathbb{Z}) \to H^k(X; \mathbb{Z})$.

Proof. The last equality is a consequence of the general theory of a spectral sequence. We prove the first two isomorphisms. For each p-cell $e^{(p)}$ of B_X, let $\tau\colon D^p \to B_X$ be the characteristic map of $e^{(p)}$ with $\tau(0) = c^{(p)}$, where $D^p (\subset \mathbb{R}^p)$ is the p-dimensional closed ball centered at 0. We put $T_X^{(p)} := \pi_{T_X}^{-1}(B_X^{(p)})$. Suppose that $e^{(p)}$ is contained in $\mathcal{S}^{(k)}B_X$. By the same way in the construction of the characteristic bundle, we can see that there exists a rank $n-k$ subtorus bundle $\pi_{S_X} : S_X \to \mathcal{S}^{(k)}B_X$ of $\pi_{T_X}|_{\mathcal{S}^{(k)}B_X} \colon T_X|_{\mathcal{S}^{(k)}B_X} \to \mathcal{S}^{(k)}B_X$ such that the restriction of $\mu_X \colon X \to B_X$ to $\mathcal{S}^{(k)}B_X$ is obtained as the quotient bundle $T_X/S_X \to \mathcal{S}^{(k)}B_X$. Since the structure group of $\pi_{T_X} : T_X \to B_X$ is discrete, there exists a bundle map $\widetilde{\tau}_T \colon (D^p, \partial D^p) \times \pi_{T_X}^{-1}(c^{(p)}) \to (T_X^{(p)}, T_X^{(p-1)})$ which covers τ such that

$$\widetilde{\tau}_T|_{\{0\} \times \pi_{T_X}^{-1}(c^{(p)})} = \mathrm{id}_{\pi_{T_X}^{-1}(c^{(p)})}$$

and $\widetilde{\tau}_T$ sends $D^p \times \pi_{S_X}^{-1}(c^{(p)})$ to S_X. Hence $\widetilde{\tau}_T$ induces a continuous map $\widetilde{\tau}_X \colon (D^p, \partial D^p) \times {\mu_X}^{-1}(c^{(p)}) \to (X^{(p)}, X^{(p-1)})$ such that

$$\widetilde{\tau}_X|_{\{0\} \times {\mu_X}^{-1}(c^{(p)})} = \mathrm{id}_{{\mu_X}^{-1}(c^{(p)})}$$

and the following diagram commutes

(8.1)
$$\begin{tikzcd}
D^p \times \pi_{T_X}^{-1}(c^{(p)}) \arrow[rrr, "\mathrm{id}_{D^p} \times \nu_{c^{(p)}}"] \arrow[dr, "\mathrm{pr}_1"] \arrow[ddr, "\widetilde{\tau}_T"'] & & & D^p \times {\mu_X}^{-1}(c^{(p)}) \arrow[dll, "\mathrm{pr}_1"'] \arrow[dd, "\widetilde{\tau}_X"] \\
& D^p \arrow[ddr, "\tau"] & & \\
& T_X \arrow[rr, "\nu"] \arrow[dr, "\pi_{T_X}"'] & & X \arrow[dl, "\mu_X"] \\
& & B_X. &
\end{tikzcd}$$

Let $\{(E_T)_r^{p,q}, d_r^T\}$ be the cohomology Serre spectral sequence of the torus bundle $\pi_{T_X} : T_X \to B_X$. By using $\widetilde{\tau}_T$, the excision isomorphism, and the Künneth formula, we have the isomorphisms

$$\begin{aligned}
(E_T)_1^{p,q} &= H^{p+q}((T_X)^{(p)}, (T_X)^{(p-1)}; \mathbb{Z}) \\
&\cong \sum_\lambda H^{p+q}(\pi_{T_X}^{-1}(\overline{e_\lambda^{(p)}}), \pi_{T_X}^{-1}(\overline{e_\lambda^{(p)}} - e_\lambda^{(p)}); \mathbb{Z}) \\
&\cong \sum_\lambda H^{p+q}((D_\lambda^p, \partial D_\lambda^p) \times \pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \\
&\cong \sum_\lambda H^p(D_\lambda^p, \partial D_\lambda^p; \mathbb{Z}) \otimes H^q(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \, (= C^p(B_X; \mathscr{H}_T^q)) \, ,
\end{aligned}$$

$$\begin{aligned}
(E_X)_1^{p,q} &= H^{p+q}(X^{(p)}, X^{(p-1)}; \mathbb{Z}) \\
&\cong \sum_\lambda H^{p+q}({\mu_X}^{-1}(\overline{e_\lambda^{(p)}}), {\mu_X}^{-1}(\overline{e_\lambda^{(p)}} - e_\lambda^{(p)}); \mathbb{Z}) \\
&\cong \sum_\lambda H^{p+q}((D_\lambda^p, \partial D_\lambda^p) \times {\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \\
&\cong \sum_\lambda H^p(D_\lambda^p, \partial D_\lambda^p; \mathbb{Z}) \otimes H^q({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}),
\end{aligned}$$

where the sum runs over all p-dimensional cells $e_\lambda^{(p)}$. By the naturality of these isomorphisms and the commutativity of the diagram (8.1), the following diagram commutes

(8.2)
$$\begin{tikzcd}
(E_X)_1^{p,q} \arrow[d, "\nu^*"'] \arrow[r, phantom, "\cong"] & \sum_\lambda H^p(D_\lambda^p, \partial D_\lambda^p; \mathbb{Z}) \otimes H^q({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \arrow[d, "\sum_\lambda \mathrm{id}_{D^p}^* \otimes \nu_{c_\lambda^{(p)}}^*"] \\
(E_T)_1^{p,q} \arrow[r, phantom, "\cong"] & \sum_\lambda H^p(D_\lambda^p, \partial D_\lambda^p; \mathbb{Z}) \otimes H^q(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}).
\end{tikzcd}$$

Moreover, it is easy to see that the homomorphism $\nu_{c_\lambda^{(p)}}^* : H^*({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \to H^*(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z})$ induced by $\nu_{c^{(p)}}$ is injective. This implies $(E_X)_1^{p,q} \cong C^p(B_X; \mathscr{H}_X^q)$. The E_1-term $\{(E_T)_1^{p,q}, d_1^T\}$ of the Serre spectral sequence is isomorphic to the CW complex $(C^p(B_X; \mathscr{H}_T^q), \delta)$ with the Serre local system $\mathscr{H}_T^q$ for the torus bundle $\pi_{T_X} : T_X \to B_X$. This fact together with the naturality of the maps in the spectral sequences shows $(E_X)_2^{p,q} \cong H^p(B_X; \mathscr{H}_X^q)$. □

Remark 8.3. (1) For $q = 0$, it is easy to see that $(E_X)_2^{p,0} \cong H^p(B_X; \mathbb{Z})$. Moreover $(E_X)_1^{p,q} = 0$, if q or p is greater than half the dimension of X.

(2) If B_X is an oriented surface with $\partial B_X \neq \emptyset$, we can take a cell decomposition of B_X so that all zero cells are included in ∂B_X. In this case, the Leray spectral sequence $\{(E_X)_r^{p,q}, d_r^X\}$ degenerates at E_2-term. In fact, $\partial B_X \neq \emptyset$ implies $(E_X)_2^{2,0} \cong H^2(B_X;\mathbb{Z}) = 0$, and since $e_\lambda^{(0)} \in \partial B_X$, the fiber ${\mu_X}^{-1}(e_\lambda^{(0)})$ of μ_X on $e_\lambda^{(0)}$ is diffeomorphic to the torus whose dimension is equal or less than one. Then $(E_X)_2^{0,2} \cong (E_X)_1^{0,2} \cong C^0(B_X;\mathscr{H}_X^2) = 0$.

Corollary 8.4. *Assume that B_X is a finite CW complex. Then the Euler characteristic $\chi(X)$ is equal to the cardinality of $\mathcal{S}^{(0)}B_X$.*

Proof. Let us consider the rational coefficient cohomology Leray spectral sequence $\{(E_X)_r^{p,q}, d_r^X\}$ of the map $\mu_X : X \to B_X$. Define

$$\chi((E_X)_r) = \sum_{p,q} (-1)^{p+q} \dim_{\mathbb{Q}} (E_X)_r^{p,q}.$$

Since $(E_X)_1^{p,q} = C^p(B_X;\mathscr{H}_X^q)$,

$$\begin{aligned}\chi((E_X)_1) &= \sum_{p,q} (-1)^{p+q} \dim_{\mathbb{Q}} (E_X)_1^{p,q} \\ &= \sum_{p,q} (-1)^{p+q} \dim_{\mathbb{Q}} C^p(B_X;\mathscr{H}_X^q) \\ &= \sum_{p,q} (-1)^{p+q} \sum_{\lambda} \dim_{\mathbb{Q}} H^q({\mu_X}^{-1}(c_\lambda^{(p)});\mathbb{Q}) \\ &= \sum_{p} \sum_{\lambda} (-1)^p \chi({\mu_X}^{-1}(c_\lambda^{(p)})),\end{aligned}$$

where the summation $\sum_\lambda$ runs over all p-cells. By the construction of $\mu_X \colon X \to B_X$, the fiber ${\mu_X}^{-1}(c_\lambda^{(p)})$ is homeomorphic to a compact torus of dimension equal or less than n, which is zero-dimensional if and only if $c_\lambda^{(p)} \in \mathcal{S}^{(0)}B_X$. Moreover, the assumption of the cell decomposition of B_X implies that $c_\lambda^{(p)} \in \mathcal{S}^{(0)}B_X$ if and only if $p = 0$. Then $\chi((E_X)_1)$ is equal to the cardinality of $\mathcal{S}^{(0)}B_X$. On the other hand, it is easy to see

$$\chi((E_X)_r) = \chi((E_X)_1)$$

for any r, and by (1) in Remark 8.3, for any r greater than n,

$$(E_X)_\infty = (E_X)_r.$$

Moreover, from $(E_X)_\infty^{p,q} = F^pH^{p+q}(X;\mathbb{Q})/F^{p+1}H^{p+q}(X;\mathbb{Q})$, we can easily check that $\chi(X) = \chi((E_X)_\infty)$. This proves the corollary. □

Example 8.5. The orbit map of Example 3.6 is equipped with a section. Let us compute the cohomology groups of X in Example 3.6. By cutting B_X along curves α and β, a cell decomposition of B_X is given. Figure 5 is the development, in which one-cells $e_1^{(1)}$, $e_2^{(1)}$, and $e_3^{(1)}$ correspond to α, β, and the edge arc γ, respectively. Let $\overline{B_X}$ be the pentagon in Figure 5. $\pi_{T_X} : T_X \to B_X$ is obtained from $\overline{B_X} \times T^2$ in the usual manner, namely, by identifying the preimage of $e_1^{(1)}$ (resp. $e_2^{(1)}$) by the first projection of $\overline{B_X} \times T^2$ with the preimage of the corresponding edge of $\overline{B_X}$ fiberwise by using $\rho([\alpha])$ (resp. $\rho([\beta])$). Let $\overline{X}$ be the quotient space

$$\overline{X} := \overline{B_X} \times T^2/\sim,$$

where $(b_1,u_1) \sim (b_2,u_2)$ if $b_1 = b_2 \in e_3^{(1)}$ and $u_2u_1^{-1} \in \{1\} \times S^1$, or $b_1 = b_2$ and it is a vertex of $\overline{B_X}$. X can be obtained from $\overline{X}$ by the same way as $\pi_{T_X} : T_X \to B_X$. The natural map $\overline{B_X} \times T^2 \to \overline{B_X} \times T^2/\sim$ descends to the map $\nu \colon T_X \to X$. For all cells except for $e^{(0)}$ and $e_3^{(1)}$, $\nu_{c_\lambda^{(p)}} : \pi_{T_X}^{-1}(c_\lambda^{(p)})(= T^2) \to {\mu_X}^{-1}(c_\lambda^{(p)})$ is a



FIGURE 5. A cell decomposition of B_X

homeomorphism, whereas for $e^{(0)}$ and $e_3^{(1)}$ it can be identified with the natural projections $T^2 \to T^2/T^2 \cong \{1\}$ and $T^2 \to T^2/\{1\} \times S^1 \cong S^1 \times \{1\}$, respectively.

For $q = 0$, $H^0(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) = \mathbb{Z}$ and $\nu^*_{c_\lambda^{(p)}} : H^0({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \to H^0(\pi_T^{-1}(c_\lambda^{(p)}); \mathbb{Z})$ is an isomorphism for all cells. Then $H^p(B_X; \mathscr{H}_X^0)$ is identified with $H^p(B_X; \mathscr{H}_T^0)$, which is naturally isomorphic to $H^p(B_X; \mathbb{Z})$.

For $q = 1$, $H^1(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) = \mathbb{Z} \oplus \mathbb{Z}$ and $\nu^*_{c_\lambda^{(p)}} : H^1({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \to H^1(\pi_T^{-1}(c_\lambda^{(p)})$ $; \mathbb{Z})$ is an isomorphism for all cells except for $e^{(0)}$ and $e_3^{(1)}$, whereas $\nu^*_{c^{(0)}}$ is the zero map and $\nu^*_{c_3^{(1)}}$ can be identified with the natural inclusion $\mathbb{Z} \oplus \{0\} \to \mathbb{Z} \oplus \mathbb{Z}$. Then $C^0(B_X; \mathscr{H}_X^1) = 0$, $C^1(B_X; \mathscr{H}_X^1) \cong \mathbb{Z}^{\oplus 5}$, and $C^2(B_X; \mathscr{H}_X^1) \cong \mathbb{Z} \oplus \mathbb{Z}$ and all coboundary operators vanish except for $\delta^1 \colon C^1(B_X; \mathscr{H}_X^1) \to C^2(B_X; \mathscr{H}_X^1)$. For $c \in C^1(B_X; \mathscr{H}_X^1)$, $\delta^1 c$ is written as

$$\begin{aligned}\delta^1 c(e^{(2)}) =& c(e_1^{(1)}) +{}^t \rho([\alpha])^{-1} c(e_2^{(1)}) -{}^t \rho([\alpha][\beta][\alpha]^{-1})^{-1} c(e_1^{(1)}) \\ & -{}^t \rho([\alpha][\beta][\alpha]^{-1}[\beta]^{-1})^{-1} c(e_2^{(1)}) +{}^t \rho([\alpha][\beta][\alpha]^{-1}[\beta]^{-1})^{-1} c(e_3^{(1)}).\end{aligned}$$

With the identification $C^1(B_X; \mathscr{H}_X^1) \cong \mathbb{Z}^{\oplus 5}$ and $C^2(B_X; \mathscr{H}_X^1) \cong \mathbb{Z} \oplus \mathbb{Z}$, δ^1 can be identified with the matrix

$$\begin{pmatrix} -1 & 1 & -2 & 2 & 3 \\ -1 & 1 & -1 & 1 & 1 \end{pmatrix}.$$

Then, the cohomology groups are calculated by

$$H^p(B_X; \mathscr{H}_X^1) = \begin{cases} \mathbb{Z}^{\oplus 3} & p = 1 \\ 0 & \text{otherwise.} \end{cases}$$

For $q = 2$, $H^2(\pi_{T_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) = \mathbb{Z}$ and $\nu^*_{c_\lambda^{(p)}} : H^2({\mu_X}^{-1}(c_\lambda^{(p)}); \mathbb{Z}) \to H^2(\pi_T^{-1}(c_\lambda^{(p)}); \mathbb{Z})$ is an isomorphism for all cells except for $e^{(0)}$ and $e_3^{(1)}$, whereas both of $\nu^*_{c^{(0)}}$ and $\nu^*_{c^{(1)}{}_3}$ are the zero map. Then $C^0(B_X; \mathscr{H}_X^2) = 0$, $C^1(B_X; \mathscr{H}_X^2) \cong \mathbb{Z}^{\oplus 2}$, and $C^2(B_X; \mathscr{H}_X^2) \cong \mathbb{Z}$. All coboundary operators vanish. It is clear except for $p = 1$. Since all monodromies along $e_\lambda^{(1)}$ induce the identity map of the second cohomology group of the fiber of $\pi_{T_X} : T_X \to B_X$ and $c(e_3^{(1)}) = 0$ for $c \in C^1(B_X; \mathscr{H}_X^2)$, δ^1 also vanishes. Then the cohomology groups are obtained by

$$H^p(B_X; \mathscr{H}_X^2) = \begin{cases} \mathbb{Z}^{\oplus 2} & p = 1 \\ \mathbb{Z} & p = 2 \\ 0 & \text{otherwise.} \end{cases}$$

The table for the E_2-terms is in Table 1. In particular, the Leray spectral sequence is degenerate at E^2-term. Then the cohomology groups of X are given by

$$H^k(X;\mathbb{Z}) = \begin{cases} \mathbb{Z} & k=0,4 \\ \mathbb{Z}^{\oplus 2} & k=1,3 \\ \mathbb{Z}^{\oplus 3} & k=2 \\ 0 & \text{otherwise.} \end{cases}$$

$$\begin{array}{c|ccc}
q & & & \\
 & 0 & \mathbb{Z}^{\oplus 2} & \mathbb{Z} \\
 & 0 & \mathbb{Z}^{\oplus 3} & 0 \\
 & \mathbb{Z} & \mathbb{Z}^{\oplus 2} & 0 \\
\hline
 & & & p
\end{array}$$

Table 1. The table of $(E_X)_2^{p,q}$-terms

8.3. **K-groups.** By replacing the cohomology functor $H^*(\,)$ by the K-functor $K^*(\,)$ in the cohomology Leray spectral sequence of the map $\mu_X : X \to B_X$, the similar method is available for computing K-groups. Such a spectral sequence is called the Atiyah-Hirzebruch spectral sequence for K-groups. We also denote this spectral sequence by the same notation $\{(E_X)_r^{p,q}, d_r^X\}$. In this subsection, the result is described without proof. But the proof is almost same as in the last subsection. For K-theory, see [2] and for the spectral sequence of K-theory, see [3]. We still assume that B_X is equipped with a structure of a CW complex so that each p-cell $e^{(p)}$ is contained in some $\mathcal{S}^{(k)}B_X$ of B_X. Let $(C^p(B_X;\mathscr{K}_T^q),\delta)$ be the cochain complex of the CW complex B_X with the local system $\mathscr{K}_T^q$ with respect to the qth K-group of the fiber of the fiber bundle $\pi_{T_X} : T_X \to B_X$. We denote by $C^p(B_X;\mathscr{K}_X^q)$ the subset of $C^p(B_X;\mathscr{K}_T^q)$ whose cochain takes a value in the image $\nu_{c^{(p)}}^*\left(K^q({\mu_X}^{-1}(c^{(p)}))\right)$ of $K^q({\mu_X}^{-1}(c^{(p)}))$ by $\nu_{c^{(p)}}^*$ for each p-cell $e^{(p)}$.

Theorem 8.6. *For any q, $C^*(B_X;\mathscr{K}_X^q)$ is a subcomplex of $(C^*(B_X;\mathscr{K}_T^q),\delta)$. We denote its pth cohomology group by $H^p(B_X;\mathscr{K}_X^q)$. Then we have the isomorphisms*

$$\begin{aligned} &(E_X)_1^{p,q} \cong C^p(B_X;\mathscr{K}_X^q), \quad (E_X)_2^{p,q} \cong H^p(B_X;\mathscr{K}_X^q), \\ &(E_X)_\infty^{p,q} = F^pK^{p+q}(X)/F^{p+1}K^{p+q}(X), \end{aligned}$$

where $F^pK^*(X)$ is the image of the map $K^*(X, X^{(p-1)}) \to K^*(X)$.

Example 8.7. Let us compute K-groups of X in Example 3.6. We also use the cell decomposition of B_X in Example 3.6. For even q, since a fiber of $\pi_{T_X} : T_X \to B_X$ is T^2, by Lemma A.1, the K-group of its fiber is isomorphic to $\mathbb{Z}\oplus\mathbb{Z}$, and all homomorphisms between them which are induced from monodromies are identity. Also by Lemma A.1, $\nu_{c_\lambda^{(p)}}^* : K^q({\mu_X}^{-1}(c_\lambda^{(p)})) \to K^q(\pi_{T_X}^{-1}(c_\lambda^{(p)}))$ is isomorphic for all cells except for $e^{(0)}$ and $e_3^{(1)}$, whereas both of $\nu_{c^{(0)}}^*$ and $\nu_{c_3^{(1)}}^*$ are the natural inclusion $\mathbb{Z}\oplus\{0\} \to \mathbb{Z}\oplus\mathbb{Z}$. A similar computation as in Example 3.6 shows that

$$H^p(B_X;\mathscr{K}_X^q) = \begin{cases} \mathbb{Z} & p=0,2 \\ \mathbb{Z}^{\oplus 4} & p=1 \\ 0 & \text{otherwise.} \end{cases}$$

For odd q, by Lemma A.3, we have an isomorphism

$$H^p(B_X; \mathscr{K}_X^q) \cong H^p(B_X; \mathscr{H}_X^1).$$

$H^p(B_X; \mathscr{H}_X^1)$ was computed in Example 3.6. The table of E^2-terms is in Table 2. In particular, the spectral sequence is degenerate at E^2-term, and K-groups of X are given by

$$\begin{array}{c|cccc} q & & & & \\ & \mathbb{Z} & \mathbb{Z}^{\oplus 4} & \mathbb{Z} & \\ & 0 & \mathbb{Z}^{\oplus 3} & 0 & \\ & \mathbb{Z} & \mathbb{Z}^{\oplus 4} & \mathbb{Z} & \\ \hline & 0 & \mathbb{Z}^{\oplus 3} & 0 & p \\ & \mathbb{Z} & \mathbb{Z}^{\oplus 4} & \mathbb{Z} & \end{array}$$

TABLE 2. The table of $(E_X)_2^{p,q}$-terms

$$K^k(X) = \begin{cases} \mathbb{Z}^{\oplus 5} & k : \text{even} \\ \mathbb{Z}^{\oplus 4} & k : \text{odd}. \end{cases}$$

8.4. **On signatures in the oriented four-dimensional case.** In this subsection, we shall give the method of computing the signature for the four-dimensional case by using the Novikov additivity. We assume that both of X and the interior of B_X are oriented so that a weakly standard atlas of X used in this subsection and the atlas of B_X induced by it are compatible with the given orientations. In this subsection the assumption $e_{orb}(X) = 0$ is not necessary.

For simplicity, suppose that B_X has only one boundary component with $\mathcal{S}^{(0)}B_X \neq \emptyset$. We divide B_X into two parts $(B_X)_1$ and $(B_X)_2$, where $(B_X)_2$ is the closed neighborhood of the boundary ∂B_X such that ∂B_X is a deformation retract of $(B_X)_2$ and $(B_X)_1$ is the closure $(B_X)_1 = \overline{B_X \backslash (B_X)_2}$ of the remainder. We put $X_i = {\mu_X}^{-1}((B_X)_i)$ for $i = 1, 2$, and denote by $\sigma(X_i)$ and $\sigma(X)$ the signature of X_i and X, respectively. The Novikov additivity says that

$$\sigma(X) = \sigma(X_1) + \sigma(X_2). \tag{8.3}$$

First let us compute $\sigma(X_1)$. As we showed in Proposition 2.7, X_1 is a T^2-bundle. If the genus of B_X is equal to zero, then $(B_X)_1$ is contractible. In this case, $\sigma(X_1)$ is zero.

Suppose that the genus of B_X is greater than zero. We give $(B_X)_1$ a trinion decomposition $(B_X)_1 = \cup_{i=1}^k ((B_X)_1)_i$, where each $((B_X)_1)_i$ is a surface obtained from S^2 by removing three distinct open discs. Let $(X_1)_i = {\mu_X}^{-1}(((B_X)_1)_i)$ for $i = 1, \dots, k$. From the Novikov additivity, we have

$$\sigma(X_1) = \sum_{i=1}^k \sigma((X_1)_i), \tag{8.4}$$

and each $\sigma((X_1)_i)$ can be computed as follows. We take oriented loops γ_1, γ_2, and γ_3 of $((B_X)_1)_i$ as in Figure 6 which represent generators of $\pi_1(((B_X)_1)_i)$ with $[\gamma_1][\gamma_2][\gamma_3] = 1$. Let $\rho : \pi_1(((B_X)_1)_i) \to Sp(2; \mathbb{Z})$ be the monodromy representation of T^2-bundle $\mu_X : (X_1)_i \to ((B_X)_1)_i$. We put $C_j := \rho([\gamma_j])$ for $j = 1, 2, 3$. For C_1 and C_2, define the vector space V_{C_1,C_2} and the bilinear form $\langle\ ,\ \rangle_{C_1,C_2}$ on V_{C_1,C_2} by

$$V_{C_1,C_2} := \{(x, y) \in \mathbb{R}^2 \times \mathbb{R}^2 : (C_1^{-1} - I)x + (C_2 - I)y = 0\},$$
$$\langle (x, y), (x', y') \rangle_{C_1,C_2} :=^t (x + y)J(I - C_2)y'$$



FIGURE 6. $((B_X)_1)_i$ and γ_j

for (x, y), $(x', y') \in V_{C_1,C_2}$, where $I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$ and $J = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. It is easy to see that $\langle\ ,\ \rangle_{C_1,C_2}$ is symmetric and we denote the signature of $\langle\ ,\ \rangle_{C_1,C_2}$ by $\tau_1(C_1, C_2)$.

Theorem 8.8 ([12, 28])**.** $\sigma((X_1)_i) = \tau_1(C_1, C_2)$.

We should remark that our orientation of $(X_1)_i$ is different from that in [12, 28]. From (8.4) and Theorem 8.8, we can compute $\sigma(X_1)$.

Remark 8.9. Meyer shows in [28] that τ_1 defines the cocycle $\tau_1 : Sp(2;\mathbb{Z}) \times Sp(2;\mathbb{Z}) \to \mathbb{Z}$ of $Sp(2;\mathbb{Z})$. It is called Meyer's signature cocycle. The author was taught Meyer's signature cocycle by Endo [12].

Next, we shall compute $\sigma(X_2)$. For X_2, we can show the following lemma.

Lemma 8.10. *${\mu_X}^{-1}(\partial B_X)$ is a deformation retract of X_2.*

Proof. Let $h_s\colon (B_X)_2 \to (B_X)_2$ $(s \in I)$ be a deformation retraction with $h_0 = \mathrm{id}_{(B_X)_2}$, $h_1((B_X)_2) = \partial B_X$, and $h_s|_{\partial B_X} = \mathrm{id}_{\partial B_X}$. By using the homotopy lifting property of the restriction of $\pi_{T_X}\colon T_X \to B_X$ to $(B_X)_2$, there exists a lift $\tilde{h}_s^T\colon T_X|_{(B_X)_2} \to T_X|_{(B_X)_2}$ of h_s such that $\tilde{h}_0^T = \mathrm{id}_{T_X|_{(B_X)_2}}$ and $\tilde{h}_s^T|_{(T_X|_{\partial B_X})} = \mathrm{id}_{T_X|_{\partial B_X}}$. $\tilde{h}_s^T$ induces the map $\tilde{h}_s^X\colon X_2 \to X_2$ such that $\tilde{h}_0^X = \mathrm{id}_{X_2}$, $\tilde{h}_s^X|_{{\mu_X}^{-1}(\partial B_X)} = \mathrm{id}_{{\mu_X}^{-1}(\partial B_X)}$, and the following diagram commutes

$$\begin{tikzcd}
 & T_X|_{(B_X)_2} \arrow[rr, "\tilde{h}_s^T"] \arrow[dl, "\nu"'] \arrow[dd, "\pi_{T_X}"] & & T_X|_{(B_X)_2} \arrow[dl, "\nu"'] \arrow[dd, "\pi_{T_X}"] \\
X_2 \arrow[rr, "\tilde{h}_s^X"] \arrow[dr, "\mu_X"'] & & X_2 \arrow[dr, "\mu_X"'] & \\
 & (B_X)_2 \arrow[rr, "h_s"] & & (B_X)_2.
\end{tikzcd}$$

In particular, $\tilde{h}_s^X$ is a required deformation retraction. $\square$

Suppose that the cardinality of $\mathcal{S}^{(0)}B_X$ is equal to k. Then $\mathcal{S}^{(1)}B_X$ has exactly k connected components $(\mathcal{S}^{(1)}B_X)_1, \ldots, (\mathcal{S}^{(1)}B_X)_k$, and ${\mu_X}^{-1}(\partial B_X) = \cup_{i=1}^k {\mu_X}^{-1}(\overline{(\mathcal{S}^{(1)}B_X)_i})$, where $\overline{(\mathcal{S}^{(1)}B_X)_i}$ is the closure of $(\mathcal{S}^{(1)}B_X)_i$. By the construction of the canonical model, it is easy to see that each ${\mu_X}^{-1}(\overline{(\mathcal{S}^{(1)}B_X)_i})$ is homeomorphic to the two-dimensional sphere S^2 if $k \geq 2$, and is homeomorphic to S^2 with one self-intersection at north and south points if $k = 1$. If $\overline{(\mathcal{S}^{(1)}B_X)_i} \cap \overline{(\mathcal{S}^{(1)}B_X)_j} \neq \emptyset$ for $i \neq j$, then they have two intersections if $k = 2$, and have one intersection if $k > 2$. In all cases, every intersection is transversal since a neighborhood of an intersection in X is identified with that of the intersection of $\mathbb{C} \times \{0\}$ and $\{0\} \times \mathbb{C}$ in $\mathbb{C}^2$. Then ${\mu_X}^{-1}(\partial B_X)$ looks like a necklace consisting of k spheres and the homology group of X_2 is given by

$$H_p(X_2;\mathbb{Z}) = H_p({\mu_X}^{-1}(\partial B_X);\mathbb{Z}) = \begin{cases} \mathbb{Z} & p = 0, 1 \\ \mathbb{Z}^{\oplus k} & p = 2 \\ 0 & \text{otherwise.} \end{cases}$$

Moreover, by putting $S_i^2 := {\mu_X}^{-1}(\overline{(\mathcal{S}^{(1)}B_X)_i})$ for $i = 1, \dots k$, the homology classes $[S_i^2] \in H_2(X_2;\mathbb{Z})$ represented by S_i^2 are generators of $H_2(X_2;\mathbb{Z})$. As a summary of the above observation, we obtain the following proposition.

Proposition 8.11. *For $i \neq j$, the intersection number $[S_i^2] \cdot [S_j^2]$ of $[S_i^2]$ and $[S_j^2]$ is given as follows*

$$[S_i^2] \cdot [S_j^2] = \begin{cases} 0 & S_i^2 \cap S_j^2 = \emptyset \\ 1 & S_i^2 \cap S_j^2 \neq \emptyset \textit{ and } k > 2 \\ 2 & S_i^2 \cap S_j^2 \neq \emptyset \textit{ and } k = 2. \end{cases}$$

Assume that $k > 1$. Let us compute the self-intersection number of $[S_i^2]$. We can take a contractible neighborhood U of $\overline{(\mathcal{S}^{(1)}B_X)_i}$ in B_X so that $U \cap \mathcal{S}^{(1)}B_X$ has exactly two connected components, say $(U \cap \mathcal{S}^{(1)}B_X)_1$ and $(U \cap \mathcal{S}^{(1)}B_X)_2$, except for $(\mathcal{S}^{(1)}B_X)_i$. We may assume that $(U \cap \mathcal{S}^{(1)}B_X)_1$ and $(U \cap \mathcal{S}^{(1)}B_X)_2$ are located as in Figure 7. Since U is contractible, there exists a local trivialization


B_X
U
$(\mathcal{S}^{(1)}(B_X))_i$
$(U \cap \mathcal{S}^{(1)}(B_X))_1$
$(U \cap \mathcal{S}^{(1)}(B_X))_2$

FIGURE 7. $(\mathcal{S}^{(1)}B_X)_i$ and $(U \cap \mathcal{S}^{(1)}B_X)_a$

$\varphi^\Lambda \colon \pi_{\Lambda_X}^{-1}(U) \cong U \times \Lambda$ of $\pi_\Lambda \colon \Lambda_X \to B_X$ which sends restrictions of $\pi_{\mathcal{L}_X} \colon \mathcal{L}_X \to \mathcal{S}^{(1)}B_X$ to $(\mathcal{S}^{(1)}B_X)_i$ and $(U \cap \mathcal{S}^{(1)}B_X)_a$ ($a = 1, 2$) fiberwise to the trivial rank one subbundles $(\mathcal{S}^{(1)}B_X)_i \times L$ and $(U \cap \mathcal{S}^{(1)}B_X)_a \times L_a$, respectively. Since $\pi_{\mathcal{L}_X} \colon \mathcal{L}_X \to \mathcal{S}^{(1)}B_X$ is primitive, we can take generators v, v_a of L and L_a such that both of the determinants of (v_1, v) and (v, v_2) are equal to one, where (v_1, v) (resp. (v, v_2)) denotes the matrix given by arranging the column vectors v_1 and v (resp. v and v_2) in this order.

Proposition 8.12. *The self-intersection number $[S_i^2] \cdot [S_i^2]$ is equal to the negative determinant $-\det(v_1, v_2)$ of (v_1, v_2).*

Note that the determinant of (v_1, v_2) does not depend on the choice of the local trivialization φ^Λ since the structure group of the bundle preserves the orientation of a fiber.

Proof. Since the self-intersection number of S_i^2 is equal to the Euler number of the normal bundle $\mathcal{N}_{S_i^2}$ of S_i^2 in X (for example, see [4]), we identify $\mathcal{N}_{S_i^2}$. For a positive number ε, let D_ε be the two-dimensional closed disc $D_\varepsilon := \{z \in \mathbb{C} \colon |z|^2 \leq \varepsilon\}$ and $\operatorname{Int} D_\varepsilon$ its interior $\operatorname{Int} D_\varepsilon := \{z \in \mathbb{C} \colon |z|^2 < \varepsilon\}$. We divide U into U_1 and U_2 as in Figure 8. We may assume that there are homeomorphisms $\varphi_1^X \colon {\mu_X}^{-1}(U_1) \to D_{\varepsilon_1} \times \operatorname{Int} D_{\varepsilon_2}$, $\varphi_2^X \colon {\mu_X}^{-1}(U_2) \to \operatorname{Int} D_{\eta_1} \times D_{\eta_2}$, $\varphi_1^B \colon U_1 \to [0, \varepsilon_1] \times [0, \varepsilon_2)$, and $\varphi_2^B \colon U_2 \to [0, \eta_1) \times [0, \eta_2]$ such that the overlap map $\varphi_{12}^X := \varphi_1^X \circ (\varphi_2^X)^{-1} \colon \operatorname{Int} D_{\eta_1} \times \partial D_{\eta_2} \to \partial D_{\varepsilon_1} \times \operatorname{Int} D_{\varepsilon_2}$ is ρ_{12}-equivariant (with respect to the standard T^2-actions) for some positive numbers ε_i, η_i ($i = 1$, 2) and $\rho_{12} \in \operatorname{Aut}(T^2)$ and overlap maps satisfy the equation $\varphi_{12}^B \circ \mu_{\mathbb{C}^2} = \mu_{\mathbb{C}^2} \circ \varphi_{12}^X$. Moreover, by applying the same argument used in Proposition 5.1 if necessary, we may also assume that overlap maps satisfy the equation $\iota \circ \varphi_{12}^B = \varphi_{12}^X \circ \iota$, where ι is the section of $\mu_{\mathbb{C}^2}$ defined by (2.2). Since φ_{12}^X sends $\{0\} \times \partial D_{\eta_2}$ to $\partial D_{\varepsilon_1} \times \{0\}$ and stabilizers of the restriction of the standard T^2-action to $\{0\} \times \partial D_{\eta_2}$ and $\partial D_{\varepsilon_1} \times \{0\}$ are $S^1 \times \{1\}$ and $\{1\} \times S^1$, respectively, ρ_{12} must be of the form



FIGURE 8. U_1 and U_2

$$\rho_{12} = \begin{pmatrix} 0 & -1 \\ 1 & m \end{pmatrix} \tag{8.5}$$

for some integer m. We show that the Euler number of the normal bundle $\mathcal{N}_{S_i^2}$ of S_i^2 in X is equal to $-m$. In this setting, S_i^2 is obtained by gluing $\{0\} \times D_{\eta_2}$ and $D_{\varepsilon_1} \times \{0\}$ with the overlap map $\varphi_{12}^X \colon \{0\} \times \partial D_{\eta_2} \to \partial D_{\varepsilon_1} \times \{0\}$. Under the natural identifications of both $\{0\} \times D_{\eta_2}$ and $D_{\varepsilon_1} \times \{0\}$ with the unit disc D_1, the direct computation using (8.5) and the fact that φ_{12}^X is ρ_{12}-equivariant shows that $\varphi_{12}^X \colon \{0\} \times \partial D_{\eta_2} \to \partial D_{\varepsilon_1} \times \{0\}$ is of the form

$$\varphi_{12}^X(z) = z^{-1}$$

for $z \in \partial D_1$ and the frame bundle of $\mathcal{N}_{S_i^2}$ can be identified with the S^1-bundle which is obtained by gluing two copies of $D_1 \times S^1$ with the map sending $(z, u) \in \partial D_1 \times S^1$ to $(z^{-1}, z^m u) \in \partial D_1 \times S^1$. This implies that the Euler number of $\mathcal{N}_{S_i^2}$ is equal to $-m$. $\square$

From Proposition 8.11 and Proposition 8.12, we can compute $\sigma(X_2)$ case-by-case for $k > 1$. In case of $k = 1$, we have a unique point b in $\mathcal{S}^{(0)} B_X$ and the fiber ${\mu_X}^{-1}(b)$ also consists of exactly one point, say $x \in {\mu_X}^{-1}(b)$. By blowing up X at x, we can reduce to the case of $k > 1$ as in the following example.

Example 8.13. Let us compute the signature of Example 3.6. Recall that B_X is a surface of genus one with one corner. As in Figure 9, we divide B_X into two parts $(B_X)_1$ and $(B_X)_2$ and give $(B_X)_1$ the trinion decomposition. The easy computation shows that the value $\tau_1(\rho([\alpha^{-1}]), \rho([\gamma^{-1}]))$ of the Meyer cocycle vanishes. This implies the signature $\sigma(X_1)$ of X_1 is zero.



the trinion decomposition of $(B_X)_1$

FIGURE 9. B_X, $(B_X)_i$, and the trinion decomposition of $(B_X)_1$

Next we focus on X_2. Since $k = 1$, the fiber of $\mu_X \colon X \to B_X$ at the unique point in $\mathcal{S}^{(0)}B_X$ consists of exactly one point which we denote by x_0. By definition, a sufficiently small neighborhood of x_0 is identified with that of the origin of $\mathbb{C}^2$. We denote the blowing-up of X_2 at x_0 by $\widetilde{X_2}$ and also denote the corresponding orbit space by $\widetilde{(B_X)}_2$. (See Example 3.15 for blowing-ups.) Note that $\mathcal{S}^{(0)}\widetilde{(B_X)}_2$ consists of exactly two points. Let $(\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_1$ and $(\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_2$ denote connected components of $\mathcal{S}^{(1)}\widetilde{(B_X)}_2$ as in Figure 10. Since the preimage $S^2_1 = {\mu_X}^{-1}((\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_1)$ is an exceptional divisor, its self-intersection number $[S^2_1] \cdot [S^2_1]$ is equal to -1. We compute the self-intersection number $[S^2_2] \cdot [S^2_2]$ of $S^2_2 = {\mu_X}^{-1}((\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_2)$. By the construction of X_2 in Example 3.6, we can take u_1, u_2 in Proposition 8.12 of the forms


$\begin{pmatrix} 0 \\ 1 \end{pmatrix}$
$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$
$(\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_1$
u_1
u_2
$(\mathcal{S}^{(1)}\widetilde{(B_X)}_2)_2$
$\begin{pmatrix} 1 \\ 0 \end{pmatrix}$
U
$(B_X)_2$
$\widetilde{(B_X)}_2$

FIGURE 10. The blowing up of $(B_X)_2$

$$u_1 = \begin{pmatrix} 3 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} 1 \\ 1 \end{pmatrix} = \begin{pmatrix} 4 \\ -1 \end{pmatrix}, \quad u_2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix},$$

hence $[S^2_2]\cdot[S^2_2] = -\det(u_1, u_2) = -5$. The above computation and Proposition 8.11 for $k = 2$ show that the intersection matrix of $\widetilde{X_2}$ is

$$\begin{pmatrix} -1 & 2 \\ 2 & -5 \end{pmatrix}$$

and $\sigma(\widetilde{X_2})$ is equal to -2. Since $\widetilde{X_2}$ is the blowing-up of X_2 at x_0, $\sigma(X_2)$ is equal to -1. Then the computations of $\sigma(X_1)$ and $\sigma(X_2)$ together with (8.3) shows that $\sigma(X) = -1$.

Appendix A. K-theory of low dimensional finite CW complexes

In this appendix, we shall give some lemmas for K-theory of low dimensional finite CW complexes which are used in Section 8.3. Let X be a finite CW complex.

Lemma A.1 ([14])**.** *If* $\dim X \leq 3$*,* $K(X)$ *is isomorphic to* $H^0(X;\mathbb{Z}) \oplus H^2(X;\mathbb{Z})$*, where* $H^0(X;\mathbb{Z}) \oplus H^2(X;\mathbb{Z})$ *is considered as a subring of the cohomology ring of* X *with* $\mathbb{Z}$*-coefficient.*

Proof. We denote $H^0(X;\mathbb{Z})\oplus H^2(X;\mathbb{Z})$ by $H^0\oplus H^2$. For simplicity we assume that X is connected. Let $\mathrm{Vect}(X)$ be the set of isomorphism classes of vector bundles on X. $\mathrm{Vect}(X)$ is a semi-ring with respect to a direct sum and a tensor product. We define the semi-ring homomorphism from $\mathrm{Vect}(X)$ to $H^0 \oplus H^2$ by assigning the rank and the first Chern class to an element of $\mathrm{Vect}(X)$. By universality of $K(X)$ it induces a ring homomorphism from $K(X)$ to $H^0 \oplus H^2$. We show that this is isomorphism. For any $(r, \alpha) \in H^0 \oplus H^2$ there exists a line bundle L on X whose first Chern class is equal to α since H^2 is identified with the set of isomorphism classes of line bundles on X. Then the above map sends the isomorphism class of $L \oplus \underline{\mathbb{C}}^{r-1}$ to (r, α), where $\underline{\mathbb{C}}^{r-1}$ is the trivial rank $(r-1)$ vector bundle. This proves that it is surjective. Next we show that it is injective. Suppose that an element $[E] - [F] \in K(X)$ lies in the kernel of this map. By construction they have the same rank, say r, and the same first Chern class. Then there exists an isomorphism $\varphi\colon \wedge^r E \to \wedge^r F$ between the top exterior bundles because they are line bundles whose first Chern classes are same as that of E and F, respectively. For $x \in X$ define Q_x by

$$Q_x := \{\text{a linear isormophism } f\colon E_x \to F_x \ :\ \wedge^r f = \varphi|_{\wedge^r E_x}\}.$$

It is easy to see that Q_x is identified with $\mathrm{SL}_r(\mathbb{C})$ and the natural projection $\pi\colon \coprod_x Q_x \to X$ is a fiber bundle with fiber $\mathrm{SL}_r(\mathbb{C})$. To prove the lemma it is sufficient to show that $\pi\colon \coprod_x Q_x \to X$ has a section. We show this by induction on the number of cells contained in X. If X consists of only one cell, it is trivial. Assume that it is valid up to $k-1$ and also assume that X consists of exactly k cells. Then X is obtained from some CW complex X' consisting of exactly $k-1$ cells by attaching a cell D. By the assumption of the induction, the restriction of $\pi\colon \coprod_x Q_x \to X$ to X' is equipped with a section, and also by the assumption of the lemma, the dimension of D is equal or less than 3. Since the ith homotopy group $\pi_i(\mathrm{SL}_r(\mathbb{C}))$ vanishes for $i \leq 2$ the section can be exteded to X. This prove the lemma. ☐

Corollary A.2. *Under the assumption of Lemma* A.1*, by assigning the first Chern class to an element of* $\widetilde{K}(X)$*, we have the ring isomorphism*

$$\widetilde{K}(X) \cong H^2(X;\mathbb{Z}). \tag{A.1}$$

Lemma A.3. *If* $\dim X \leq 2$*, there is a ring isomorphism*

$$K^{-1}(X) \cong H^1(X;\mathbb{Z})$$

which commutes with maps induced from a continuous map $f : X \to Y$

$$\begin{tikzcd}
K^{-1}(X) \arrow[r, "\cong"] & H^1(X;\mathbb{Z}) \\
K^{-1}(Y) \arrow[r, "\cong"] \arrow[u, "f^*"] \arrow[ur, phantom, "\circlearrowright"] & H^1(Y;\mathbb{Z}). \arrow[u, "f^*"']
\end{tikzcd}$$

Proof. By definition,

$$K^{-1}(X) := \widetilde{K}(\widetilde{S}X^+),$$

where X^+ is the disjoint union of X with a point x_0 and $\widetilde{S}X^+$ is the reduced suspension of X^+. Let SX^+ and Sx_0 be unreduced suspensions of X^+ and x_0, respectively. Sx_0 is naturally included in SX^+ and the quotient space obtained from SX^+ by collapsing Sx_0 is just $\widetilde{S}X^+$. Atiyah shows in [2] that the natural projection $p : SX^+ \to \widetilde{S}X^+$ induces the isomorphism

$$p^* : \widetilde{K}(\widetilde{S}X^+) \cong \widetilde{K}(SX^+). \tag{A.2}$$

By composing the isomorphisms (A.2), (A.1) for SX^+, and the suspension isomorphism

$$H^2(SX^+;\mathbb{Z}) \cong H^1(X^+;\mathbb{Z})(\cong H^1(X;\mathbb{Z})),$$

we obtain the isomorphism in the lemma. □

Remark A.4. It is also shown in [14] that when $\dim X \le 5$, by associating the rank, the first Chern class, and the second Chern class with an element of $K(X)$ (by associating the first and second Chern classes in the case of $\widetilde{K}(X)$), we have the one-to-one correspondences as a set

$$K(X) \cong H^0(X;\mathbb{Z}) \oplus H^2(X;\mathbb{Z}) \oplus H^4(X;\mathbb{Z}),$$
$$\widetilde{K}(X) \cong H^2(X;\mathbb{Z}) \oplus H^4(X;\mathbb{Z}).$$

Then the same argument in the proof of Lemma A.3 shows that when $\dim X \le 4$, there is an one-to-one correspondence

$$K^{-1}(X) \cong H^1(X;\mathbb{Z}) \oplus H^3(X;\mathbb{Z})$$

as a set.

References

1. V. I. Arnold, *Mathematical methods of classical mechanics*, second ed., Graduate texts in Mathematics, vol. 60, Springer-Verlag, New York, 1989.
2. M. F. Atiyah, *K-theory*, Benjamin, New York, 1967.
3. M. F. Atiyah and F. Hirzebruch, *Vector bundles and homogeneous spaces*, Proc. Sympos. Pure Math. (Providence, RI), vol. III, Amer. Math. Soc., 1961, pp. 7–38.
4. R. Bott and L. W. Tu, *Differential forms in algebraic topology*, Graduate texts in Mathematics, vol. 82, Springer-Verlag, New York-Berlin, 1982.
5. M. Boucetta and Molino P., *Géométrie globale des systèmes hamiltoniens complètement intègrables : fibrations lagrangiennes singulières et coordonnées action-angle à singularités*, C. R. Acad. Sci. Paris Sér. I Math. **308** (1989), no. 13, 421–424.
6. V. Buchstaber and T. Panov, *Torus actions and their applications in topology and combinatorics*, University Lecture Series, vol. 24, Amer. Math. Soc., Providence, RI, 2002.
7. V. Buchstaber and N. Ray, *Tangential structures on toric manifolds and connected sums of polytopes*, Internat. Math. Res. Notices (2001), no. 4, 193–219.
8. V. Danilov, *The geometry of toric varieties (Russian)*, Uspekhi Mat. Nauk **33** (1978), no. 2, 85–134, English translation: Russian Math. Surveys 33 (1978), no. 2, 97–154.
9. M. Davis, *Group generated by reflections and aspherical manifolds not covered by Euclidean space*, Ann. of Math. (2) **117** (1983), no. 2, 293–324.
10. M. Davis and T. Januszkiewicz, *Convex polytopes, coxeter orbifolds and torus actions*, Duke Math. J. **62** (1991), no. 2, 417–451.

11. J. J. Duistermaat, *On global action-angle coordinates*, Comm. Pure Appl. Math. **33** (1980), no. 6, 687–706.
12. H. Endo, *Lefschetz fibrations and Meyer's signature cocycle*, Nov. 2005, The intensive lecture given at Graduate School of Mathematical Sciences, The University of Tokyo.
13. W. Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993.
14. M. Furuta, *Topological K-theory*, 2003, The lecture given at Graduate School of Mathematical Sciences, The University of Tokyo.
15. D. T. Gay and M. Symington, *Toric structures on near-symplectic* 4*-manifolds*, arXiv:math.SG/0609753, 2006.
16. H. Geiges, *Symplectic structures on T^2-bundles over T^2*, Duke Math. J. **67** (1992), no. 3, 539–555.
17. V. Guillemin and S. Sternberg, *Birational equivalence in the symplectic category*, Invent. Math. **97** (1989), no. 3, 485–522.
18. A. Haefliger and É. Salem, *Actions of tori on orbifolds*, Ann. Global Anal. Geom. **9** (1991), no. 1, 37–59.
19. M. Hamilton, *Locally toric manifolds and singular Bohr-Sommerfeld leaves*, arXiv:math.SG/0709.4058, 2007.
20. A. Hattori, *Topology I–III*, second ed., Iwanami Lectures on Fundamental Mathematics, vol. Geometry ii, Iwanami Shoten, Tokyo, 1982, English translation: Isōkikagaku I–III. (Japanese), Iwanami Kōza Kiso Sūgaku, Kikagaku ii, Iwanami Shoten, Tokyo, 1982.
21. ______, *Elliptic genera, torus orbifolds and multi-fans. II*, Internat. J. of Math. **17** (2006), no. 6, 707–735.
22. A. Hattori and M. Masuda, *Theory of multi-fans*, Osaka J. Math. **40** (2003), no. 1, 1–68.
23. ______, *Elliptic genera, torus manifolds and multi-fans*, Internat. J. of Math. **16** (2005), no. 9, 957–998.
24. H. B. Lawson and M. L. Michelsohn, *Spin geometry*, Princeton Mathematical Series, vol. 38, Princeton University Press, Princeton, NJ, 1989.
25. E. Lerman, *Symplectic cuts*, Math. Res. Lett. **2** (1995), no. 3, 247–258.
26. M. Masuda, *Unitary toric manifolds, multi-fans, and equivariant index*, Tohoku Math. J. **51** (1999), no. 2, 237–265.
27. M. Masuda and T. Panov, *On the cohomology of torus manifolds*, Osaka J. Math. **43** (2006), no. 3, 711–746.
28. W. Meyer, *Die signaturvon flächenbündeln*, Math. Ann. **201** (1973), no. 3, 239–264.
29. K. N. Mishachev, *The classification of lagrangian bundles over surface*, Differential Geom. Appl. **6** (1996), no. 4, 301–320.
30. T. Oda, *Convex bodies and algebraic geometry. an introduction to the theory of toric varieties*, Springer-Verlag, Berlin, 1988, Translated from Japanese.
31. P. Orlik and F. Raymond, *Actions of the torus on* 4*-manifolds. I*, Trans. Amer. Math. Soc. **152** (1970), no. 2, 531–559.
32. M. Symington, *Four dimensions from two in symplectic topology*, Topology and geometry of manifolds (Athens, GA, 2001), Proc. Sympos. Pure Math., vol. 71, Amer. Math. Soc., Providence, RI, 2003, pp. 153–208.
33. T. Yoshida, *Twisted toric structures*, arXiv:math.SG/0605376, 2006.
34. ______, *On liftings of local torus actions to fiber bundles*, to appear in the Proceedings of International Conference on Toric Topology, 2007.

Graduate School of Mathematical Sciences, The University of Tokyo,
8-1 Komaba 3-chome, Meguro-ku, Tokyo, 153-8914, Japan
E-mail address: `takahiko@ms.u-tokyo.ac.jp`